

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2014 ^{1, 2, 3}

March 23, 2016 download of January 2014 – December 2014 NHSN data

Hospitals	Type of Surgical Procedure ⁴																																
	Abdominal Hysterectomy			Colon Surgery			Coronary Artery Bypass Graft ⁵						Heart Surgery			Hip Replacement ⁵						Knee Replacement ⁵											
							Chest and Secondary Incision Site		Chest Incision Only		Overall					Initial		Revision		Overall		Initial		Revision		Overall							
	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate				
Kadlec Medical Center	1.19	1.19	84	0.00	0.83	120	1.32	1.32	302	0.00	0.00	11	1.28	1.28	0.00	0.00	144	0.00	0.00	166	0.00	0.00	18	0.00	0.00	0.00	0.00	191	0.00	0.00	8	0.00	0.00
Kittitas Valley Healthcare ²⁹	0.00	0.00	4	0.00	0.00	5												3.39	6.78	59	0.00	0.00	8	2.99	5.97	0.00	0.00	104	0.00	0.00	8	0.00	0.00
Lake Chelan Community Hospital	0.00	0.00	1	-	-	0																											
Legacy Salmon Creek Hospital	1.53	2.29	131	3.97	5.56	126												2.16	2.16	231	0.00	0.00	19	2.00	2.00	0.31	0.31	638	0.00	0.00	45	0.29	0.29
Lincoln Hospital	0.00	0.00	7	0.00	0.00	9																											
Lourdes Health Network ¹¹	3.13	3.13	32	5.88	5.88	17												0.00	0.00	118	3.70	3.70	27	0.69	0.69	0.76	1.01	395	0.00	0.00	29	0.71	0.94
Mary Bridge Children's Hospital				0.00	0.00	23								0.00	0.00	36																	
Mason General Hospital	0.00	0.00	7	0.00	8.33	12												0.00	0.00	24	0.00	0.00	5	0.00	0.00	0.00	0.00	54	0.00	0.00	6	0.00	0.00
Mid-Valley Hospital ³⁰	0.00	0.00	8	9.09	9.09	11												4.76	4.76	21	0.00	100.00	1	4.55	9.09	0.00	0.00	24	-	-	0	0.00	0.00
Morton General Hospital ¹⁹	-	-	0	20.00	20.00	5																											
MultiCare Auburn Medical Center	1.45	1.45	69	4.92	4.92	61												0.00	0.00	91	0.00	0.00	14	0.00	0.00	0.00	0.00	143	0.00	0.00	22	0.00	0.00
MultiCare Good Samaritan Hospital	1.57	1.57	127	2.68	2.68	149												0.45	0.45	220	2.94	2.94	34	0.79	0.79	0.00	0.00	262	0.00	0.00	14	0.00	0.00
Newport Community Hospital				-	-	0																											
North Valley Hospital	-	-	0																														

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2014^{1, 2, 3}

March 23, 2016 download of January 2014 – December 2014 NHSN data

Hospitals	Type of Surgical Procedure ⁴																																	
	Abdominal Hysterectomy			Colon Surgery			Coronary Artery Bypass Graft ⁵						Heart Surgery			Hip Replacement ⁵						Knee Replacement ⁵												
							Chest and Secondary Incision Site			Chest Incision Only						Overall			Initial		Revision		Overall		Initial		Revision		Overall					
	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate					
Northwest Hospital and Medical Center – UW Medicine	0.00	0.00	47	0.91	4.55	110	0.00	0.00	37	0.00	0.00	4	0.00	0.00	0.00	0.00	21	0.00	0.57	351	0.00	0.00	42	0.00	0.51	0.00	0.26	380	0.00	0.00	28	0.00	0.25	
Ocean Beach Hospital	0.00	0.00	1	0.00	0.00	12																												
Olympic Medical Center	0.00	0.00	41	2.90	5.80	69												0.00	0.00	123	0.00	0.00	1	0.00	0.00	0.00	0.00	156	-	-	0	0.00	0.00	
Othello Community Hospital	0.00	0.00	1																															
Overlake Hospital Medical Center	0.78	1.29	387	2.94	4.58	306	0.00	0.75	133	0.00	0.00	4	0.00	0.73	0.00	0.00	93	1.11	1.11	719	0.00	0.00	69	1.02	1.02	0.12	0.12	853	0.00	0.00	59	0.11	0.11	
PeaceHealth Saint John Medical Center	0.00	0.00	42	2.50	5.00	80												0.52	0.52	193	0.00	0.00	12	0.49	0.49	0.31	0.31	320	5.88	5.88	17	0.59	0.59	
PeaceHealth Saint Joseph Medical Center	0.00	0.96	104	2.70	4.05	148	0.81	1.61	124	0.00	0.00	6	0.77	1.54	0.00	0.00	134	0.76	0.76	397	0.00	0.00	18	0.72	0.72	0.23	0.45	444	0.00	5.26	38	0.21	0.83	
PeaceHealth Southwest Medical Center	0.00	0.00	294	3.24	5.09	216	0.00	0.00	197	0.00	0.00	3	0.00	0.00	0.00	0.00	110	0.23	0.69	435	0.00	0.00	25	0.22	0.65	0.46	0.46	432	0.00	0.00	31	0.43	0.43	
PeaceHealth United General Hospital				0.00	0.00	4																												
Prosser Memorial Hospital Medical Center	0.00	0.00	7	-	-	0												0.00	0.00	7	-	-	0	0.00	0.00	0.00	0.00	15	-	-	0	0.00	0.00	
Providence Centralia Hospital ¹²	2.04	6.12	49	0.00	0.00	46												0.88	0.88	114	0.00	0.00	1	0.87	0.87	0.00	0.00	164	0.00	0.00	13	0.00	0.00	
Providence Holy Family Hospital ³¹	0.00	0.00	64	2.33	6.98	86												0.85	0.85	236*	12.00	14.00	50*	2.80	3.15*	0.00	0.00	367	1.89	1.89	53	0.24	0.24	
Providence Mount Carmel Hospital	0.00	0.00	1	0.00	0.00	10												0.00	0.00	1	-	-	0	0.00	0.00	0.00	0.00	1	-	-	0	0.00	0.00	
Providence Regional Medical Center Everett	1.21	1.93	414	1.26	2.93	239	0.98	1.64	305	0.00	0.00	10	0.95	1.59	0.00	0.66	152	0.18	0.71	564	2.56	2.56	39	0.33	0.83	0.36	0.53	563	2.70	8.11	37	0.50	1.00	

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2014^{1, 2, 3}

March 23, 2016 download of January 2014 – December 2014 NHSN data

Hospitals	Type of Surgical Procedure ⁴																																	
	Abdominal Hysterectomy			Colon Surgery			Coronary Artery Bypass Graft ⁵						Heart Surgery			Hip Replacement ⁵						Knee Replacement ⁵												
							Chest and Secondary Incision Site			Chest Incision Only						Overall			Initial		Revision		Overall		Initial		Revision		Overall					
	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate	No. of Procedures	Deep Rate	Total Rate					
Providence Sacred Heart Medical Center ²⁴	0.27	0.81	372	3.32	3.99	301	0.25	0.25	395	0.00	0.00	18	0.24	0.24	0.37	0.37	534	0.15	0.59	674	1.87	1.87	107	0.38	0.77	0.28	0.42	711	3.31	3.31	121	0.72	0.84	
Providence Saint Mary Medical Center ³³	0.00	0.00	18	6.38	8.51	47												1.42	1.42	141	0.00	0.00	13	1.30	1.30	0.56	0.56	178	5.26	5.26	19	1.02	1.02	
Providence Saint Peter Hospital	0.00	0.00	90	0.47	0.93	215	0.97	1.45	207	0.00	0.00	17	0.89	1.34	0.00	0.00	134	0.22	0.22	449	1.79	1.79	56	0.40	0.40	0.12	0.49	818	0.00	1.46	137	0.10	0.63	
Pullman Regional Hospital	0.00	1.23	81	6.25	12.50	16												0.00	0.00	41	-	-	0	0.00	0.00	0.74	1.48	135	0.00	0.00	7	0.70	1.41	
Saint Anthony Hospital ¹³	0.00	5.00	20	2.82	7.04	71												3.33	3.33	30	0.00	0.45	223	0.40	0.79	0.35	0.35	285	0.00	0.00	19	0.33	0.33	
Saint Clare Hospital ³⁴	0.00	0.00	1	0.00	0.00	45												0.00	0.00	33	0.42	1.27	236	0.37	1.12	0.74	2.21	271	0.00	0.00	19	0.69	2.07	
Saint Elizabeth Hospital	0.00	0.00	9	4.55	4.55	22												0.00	0.00	43	0.00	0.00	4	0.00	0.00	0.00	0.00	37	0.00	0.00	1	0.00	0.00	
Saint Francis Hospital	0.00	0.00	15	0.00	4.29	70												0.00	0.00	24	0.00	0.00	45	0.00	0.00	0.00	0.00	122	0.00	0.00	5	0.00	0.00	
Saint Joseph Medical Center	0.65	0.65	154	2.23	3.57	224	0.36	0.36	276	0.00	0.00	17	0.34	0.34	0.00	0.00	250	0.00	0.00	37	1.17	1.46	343	1.05	1.32	0.41	0.61	490	2.27	2.27	44	0.56	0.75	
Samaritan Healthcare	0.00	0.00	31	0.00	10.00	30												0.00	0.00	48	-	-	0	0.00	0.00	0.00	0.00	148	0.00	0.00	1	0.00	0.00	
Seattle Children's Hospital ^{25, 35}	0.00	0.00	4	1.54	4.62	65	-	-	0	-	-	0	-	-	0.61	0.61	330	0.00	0.00	1	-	-	0	0.00	0.00	25.00	25.00	4	0.00	0.00	1	20.00	20.00	
Shriners Hospital for Children Spokane																																		
Skagit Valley Hospital ³⁶	0.00	0.00	35	3.77	7.55	53												0.00	1.80	111	0.00	0.00	5	0.00	1.72	0.60	0.60	167	5.56	5.56	18	1.08	1.08	
Skyline Hospital	-	-	0	-	-	0																												

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2014^{1, 2, 3}

March 23, 2016 download of January 2014 – December 2014 NHSN data

The annual number of procedures performed by a hospital is also provided as context to explain infection rates, and inform patients and providers in planning for elective surgery. In general, surgical teams which perform more of the same procedures have better outcomes.

2. Depth of infection classification:
 - a. The deep infection rate includes surgical wound infections occurring well below the skin and subcutaneous tissue, affecting muscles, organs, or bones (“deep incisional” and “organ/space” SSI per NHSN). Deep infections may be severe and can be difficult to treat often requiring antibiotics and a return visit to a hospital.
 - b. The total infection rate includes deep infections as well as shallow surgical wound infections occurring within the skin and subcutaneous tissue (“superficial incisional” SSI per NHSN). Shallow infections can be more easily treated, often without hospitalization and may not require antibiotics. More information about the depth of [surgical site infections](#).
 - c. Note – Rates for deep and total infections from the same procedure type use the same denominator (number of procedures performed) during the reporting period. See [box plots](#) for a graphical representation of the distribution of infection rates.
3. Standardized infection ratios (SIR) are summary statistics used by CDC/NHSN, states, Centers for Medicare and Medicaid Services (CMS), and healthcare facilities to track healthcare associated infection prevention progress over time. The SIR compares reported HAI cases in a facility to “predicted” cases from historic baseline rates from the general U.S. population (national baseline for 2014 SSI SIR is January 2006 – December 2008). The SIR for a facility is adjusted to account for factors that might cause infection rates to be higher or lower, such as hospital, surgery and patient characteristics. Washington State hospitals with surgical site infection SIRs significantly different from the national baseline are identified as statistically high (red with an asterisk “*”), with low (gold/yellow) SIRs to be indicated in future reports after hospital SSI reporting is validated. Washington State reports primarily use the “All SSI SIR Model” to support outlier SSI identification.
4. Visit our website for descriptions of these [types of surgical procedure categories](#).
5. Details of certain specific surgical procedure types are included in this table (coronary artery bypass graft procedures by incision sites and initial vs. revision joint replacement procedures). Hospitals are compared by their “overall” infection rates in these categories to compare Washington hospitals more accurately. However, by separating rates into more specific procedure types, the number of procedures performed in either subcategory can be very low, and a single infection may produce a seemingly high rate. The more specific infection rate and procedure volume information in this table may be useful for patients and providers in planning for elective surgery.

Hospital Notes:

6. Trios Health / Southridge Hospitals – Most inpatient surgical hospital services (including surgeries) at Trios Health were largely discontinued mid-year in 2014 and transferred to the new Trios Southridge facility, which opened June 2014. These reported rates are the combination of both facilities’ respective surgical type rates over the year. See additional comment (#14) under Abdominal Hysterectomies.

Abdominal Hysterectomies

Infection rates and number of procedures performed for this type of surgery include only inpatient hysterectomies with abdominal incisions, per NHSN surveillance definitions. Hysterectomies where organs are detached via vaginal approach, and outpatient (procedure and hospital discharge in the same day) abdominal hysterectomies are considered different types of surgical procedures with differing risk for infection, and are not included in state surveillance for

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2014^{1, 2, 3}

March 23, 2016 download of January 2014 – December 2014 NHSN data

this abdominal hysterectomy category. Therefore, the volume of procedures reported here may not represent a hospital's overall experience performing "hysterectomies" in general.

7. Cascade Valley Hospital – Infection rate is due to one deep surgical wound infection in the only inpatient abdominal hysterectomy performed during the year. Approximately 40 hysterectomies (both abdominal and vaginal) are performed annually and most are outpatient surgeries. The hospital investigated the case and identified a gap in the preoperative skin preparation for this patient. Subsequently, the hospital reports staff reeducation on surgical care measures.
8. Central Washington Hospital – Infection rate is due to two surgical wound infections, one shallow and one deep, with a very low number of inpatient abdominal hysterectomies performed during the year. The hospital investigated both cases and reports surgical care measures were followed.
9. Deaconess Hospital – Infection rate is due to three surgical wound infections, one shallow and two deep, with no indication of cross-infection. The hospital investigated the cases and reports surgical care measures were followed. Subsequently, the hospital reports increasing surveillance among obstetrics-gynecology and oncology surgery patients to find additional cases; no trends were identified.
10. Harborview Medical Center – Infection rate is due to two shallow surgical wound infections, with no indication of cross-infection. The hospital investigated both cases and reports that surgical care and infection prevention measures were followed. The hospital continues to report enhanced infection surveillance due to a large outpatient clinic network.
11. Lourdes Health Network - Infection rate is due to only one deep surgical wound infection. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed.
12. Providence Centralia Hospital – Infection rate is due to three surgical wound infections, two shallow and one deep. The hospital investigated each case and identified variable practices with preoperative skin preparation and environmental cleaning. Subsequently, the hospital has added training and competency assessments for preoperative skin preparation; enhanced patient focused preoperative infection prevention measures; and re-evaluated environmental cleaning products and processes.
13. Saint Anthony Hospital – Infection rate is due to only one shallow surgical wound infection and a low number of inpatient abdominal hysterectomies performed during the year. The hospital investigated the case and reports surgical care measures were followed. Subsequent to this infection and other 2014 surgical infections across this hospital network (CHI Franciscan Health), the hospital reports various practices have changed in this facility (see footnote #34 for Saint Clare Hospital and knee replacement infections, regarding hospital system-wide interventions).
14. Trios Health / Southridge Hospitals – Infection rate is due to one deep surgical wound infection during the second half of the year (at Trios Southridge Hospital). The hospital investigated the case and reports surgical care measures were followed. Most inpatient surgical hospital services (including non-emergent surgeries) at Trios Health were largely discontinued mid-year in 2014 and transferred to the new Trios Southridge facility, which opened June 2014. This reported rate is the combination of both facilities' abdominal hysterectomy rates over the year.
15. Valley Medical Center – Infection rate is due to two deep surgical wound infections, with no indication of cross infection. The hospital investigated both cases and reports that surgical care measures were followed.
16. Whidbey General Hospital – Infection rate is due to one deep surgical wound infection with a very low number of inpatient abdominal hysterectomies performed during the year. Approximately 30 inpatient and outpatient hysterectomies (both abdominal and vaginal) are performed annually. The hospital's standardized infection ratio (SIR) for surgical site infection across each reported category combined is also significantly higher than the national baseline. The hospital investigated each case and reports that surgical care measures were followed. Subsequently, the hospital reports changes in environmental infection controls in the operating room.

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2014^{1, 2, 3}

March 23, 2016 download of January 2014 – December 2014 NHSN data

Colon Surgeries

17. Central Washington Hospital – The hospital's standardized infection ratio (SIR) for colon surgeries, as calculated by CDC/NHSN, is significantly higher than the national baseline - with two times the predicted number of infections occurring throughout the year. Of the fifteen reported surgical wound infections, the SIR includes fourteen, seven shallow and seven deep – and excludes one infection per NHSN exclusion criteria. The hospital reviewed each case and reports surgical care measures were met, excluding emergent cases, where not all routine preoperative measures could be performed. The hospital also continues to report enhanced infection surveillance from access to an outpatient clinic network.
18. Harborview Medical Center – Harborview Medical Center – The hospital's standardized infection ratio (SIR) for colon surgeries, as calculated by CDC/NHSN is significantly higher than the national baseline. Of the twenty-one reported surgical wound infections, the SIR includes eighteen, seven shallow and eleven deep – and excludes three infections per NHSN exclusion criteria. The hospital investigated each case and reports surgical care measures were followed except for multiple emergent cases, where all routine preoperative measures could not be performed. Patients receiving colon surgeries in emergent settings or for traumatic injuries may be at greater risk for developing infections – however these elements are not accounted for in risk adjusting the SIR for colon surgery. The hospital reports enhanced patient focused infection prevention measures were implemented in 2015.
19. Morton General Hospital - Infection rate is due to one deep surgical wound infection among very few colon procedures performed during the year. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed. Subsequently, enhanced patient focused infection prevention measures were implemented.
20. Swedish First Hill – The hospital's standardized infection ratio (SIR) for colon surgeries, as calculated by CDC/NHSN is significantly higher than the national baseline – with one-and-a-half times the predicted number of infections occurring throughout the year. Of the forty-two reported surgical wound infection, the SIR includes thirty-nine, fifteen shallow and twenty-four deep – and excludes three deep infections per NHSN exclusion criteria. The hospital investigated each case with a multidisciplinary team and reports surgical care measures were followed. Subsequently the hospital initiated multiple colon infection prevention activities including reinforcing surgical care measures and prophylactic antibiotic timing, and implementing enhanced environmental and patient-focused perioperative measures.
21. Swedish Issaquah – Infection rate is due to nine surgical wound infections, two shallow and seven deep, throughout the year. The hospital's standardized infection ratio (SIR) for colon surgery infection is also significantly higher than the national baseline. The hospital reports surgical care measures were followed. Subsequently the hospital initiated multiple colon infection prevention activities including reinforcing surgical care measures and prophylactic antibiotic timing, and continues to reinforce enhanced patient-focused prevention measures.
22. Whidbey General Hospital – Infection rate is due to two deep surgical wound infections, with no indication of cross-infection among the few colon procedures performed during the year. The hospital's standardized infection ratio (SIR) for surgical site infection across each reported category combined is also significantly higher than the national baseline. The hospital investigated each case and identified some gaps in surgical care measures. Subsequently, the hospital reports changes in environmental infection controls in the operating room.

Coronary Artery Bypass Graft Surgeries

For coronary artery bypass graft surgeries, hospitals are compared by their “overall” infection rates, to compare hospitals more accurately. Details of coronary artery bypass graft surgeries with secondary incision sites (CBGB) or chest incision only (CBGC) may be useful for patients and providers in planning for elective surgery and for comparison to other reports using these NHSN categories.

No Washington State hospital had statistically higher rates of coronary artery bypass graft associated infection by statewide rate comparisons or SIR for 2014.

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2014^{1, 2, 3}

March 23, 2016 download of January 2014 – December 2014 NHSN data

Heart (Cardiac) Surgeries

23. Harborview Medical Center – Infection rate is due to only one superficial surgical wound infection among very few cardiac surgeries performed during the year. The hospital investigated the case and reports that surgical care and infection prevention measures were followed. The hospital does not perform scheduled heart surgery; the patients that receive these procedures are admitted for emergent and traumatic injuries.
24. Providence Sacred Heart Medical Center – Infection rate is due to two deep surgical wound infections during the year, with no indication of cross-infection. However, the hospital's cardiac surgery standardized infection ratio (SIR) is significantly lower than the national baseline – indicating fewer reported annual infections than predicted by CDC/NHSN. The hospital investigated both cases and reports that surgical care measures were followed. Subsequently, the hospital reports implementing enhanced patient-focused and surgical care infection prevention measures.
25. Seattle Children's Hospital – Infection rate is due to two deep surgical wound infections throughout the year, with no indication of cross-infection. The hospital's cardiac surgery standardized infection ratio (SIR), which excludes one infection per NHSN exclusion criteria, is significantly lower than the national baseline – indicating fewer reported annual infections than predicted by CDC/NHSN. The hospital operates on infants and children with multiple complex anatomical anomalies undergoing reparative heart surgeries. The hospital investigated both cases and reports that surgical care and infection control measures were followed.
26. Swedish Cherry Hill – Infection rate is due to two surgical wound infections, one shallow and one deep, with no indication of cross-infection. The hospital's infection control and quality programs investigated the cases and reports complete surgical care measures were followed for one case; the one superficial case was emergent where all routine preoperative measures could not be performed. The hospital reports enhanced patient-focused and surgical-care infection prevention measures continue to be maintained.
27. Virginia Mason Medical Center – Infection rate is due to only one deep surgical wound infection. The hospital investigated the case and subsequently reports assessing operating room environmental factors and implementing enhanced patient-focused infection prevention measures.

Hip Replacement Surgeries

For hip replacement surgeries, hospitals are compared by their "overall" infection rates, to compare hospitals more accurately. Details of initial ("primary" per NHSN) and repeat, revision and resurfacing hip prosthesis surgeries may be useful for patients and providers in planning for elective surgery.

28. Harborview Medical Center – Infection rate is due to only one deep surgical wound infection among a low number of hip replacements performed during the year. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed. The hospital does not perform elective joint replacement surgery; the patients that receive these procedures are admitted for emergent and traumatic injuries.
29. Kittitas Valley Community Hospital – Infection rate is due to four surgical wound infections, two shallow and two deep. The hospital investigated each case and reports surgical care infection prevention measures were followed. Subsequently, the hospital reports enhanced patient focused infection prevention measures were implemented for patients transferred from the emergency department to the operating room, and for post-operative wound dressing.
30. Mid-Valley Hospital – Infection rate is due to two surgical wound infections, one shallow and one deep, with no indication of cross-infection and a low number of hip replacements performed during the year. The hospital investigated each case and reports that surgical care measures were followed.
31. Providence Holy Family Hospital – Infection rate is due to nine surgical wound infections throughout the year, one shallow and eight deep. The hospital's standardized infection ratio (SIR) for hip replacement infections is also significantly higher than the national baseline. The hospital investigated each case and reports that surgical care measures were followed. Seven of these cases were revisions to the original hip surgery, some necessitated by patient-falls after

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2014^{1, 2, 3}

March 23, 2016 download of January 2014 – December 2014 NHSN data

discharge. Subsequent to these cases, the hospital reports implementing enhanced patient-focused preoperative measures, reinforcing best practices for operating room environmental infection prevention, and providing surgical staff reeducation.

32. Whidbey General Hospital – Infection rate is due to two surgical wound infections, one shallow and one deep, with no indication of cross-infection and a low number of hip replacements performed during the year. The hospital's standardized infection ratio (SIR) for surgical site infection across each reported category combined is also significantly higher than the national baseline. The hospital investigated each case and reports that surgical care measures were followed. Subsequently, the hospital reports changes in environmental infection controls in the operating room and enhanced patient-focused preoperative infection prevention measures for joint replacement procedures.

Knee Replacement Surgeries

For knee replacement surgeries, hospitals are compared by their "overall" infection rates, to compare hospitals more accurately. Details of initial ("primary" per NHSN) and repeat, revision and resurfacing knee prosthesis surgeries may be useful for patients and providers in planning for elective surgery.

33. Providence Saint Mary Medical Center – Infection rate is due to two deep surgical wound infections, with no indication of cross-infection. The hospital reviewed each case and reports surgical care measures were followed. Subsequently, the hospital reports enhanced patient-focused infection prevention measures were implemented for preoperative care.
34. Saint Clare Hospital – Infection rate is due to six surgical wound infections during the year, four shallow and two deep, with no indication of cross-infection. The hospital investigated each case and reports surgical care measures were followed. Subsequent to these infections and other 2014 surgical infections across this hospital network (CHI Franciscan Health), the hospital reports various practices have changed in this facility: operative staff reeducation on surgical care measures with increased adherence monitoring; enhanced patient-focused preoperative infection prevention measures; standardizing environmental infection controls in operating rooms; and development of a multi-disciplinary group to review and investigate surgical practices and infections.
35. Seattle Children's Hospital – Infection rate is due to only one deep surgical wound infection, among very few knee replacements performed during the year. The hospital does not perform elective joint replacement surgery. The patients that receive these procedures have malignancies or complex anatomical anomalies; this patient was severely immunocompromised. The hospital investigated the case and reports surgical care and infection prevention measures were followed.
36. Skagit Valley Hospital – Infection rate is due to two deep surgical wound infections, with no indication of cross-infection. The hospital investigated each case and reports surgical care measures were followed. Subsequently, the hospital reports enhanced patient-focused infection measures were implemented for preoperative care.
37. Whidbey General Hospital – Infection rate is due to two surgical wound infections, one shallow and one deep, with no indication of cross-infection. The hospital's standardized infection ratio (SIR) for surgical site infection across each reported category combined is also significantly higher than the national baseline. The hospital investigated each case and reports that surgical care measures were followed. Subsequently, the hospital reports changes in environmental infection controls in the operating room and enhanced patient-focused preoperative infection prevention measures for joint replacement procedures.
38. Yakima Regional Medical and Cardiac Center – Infection rate is due to three deep surgical wound infections, with no indication of cross-infection. The hospital reviewed each case and reports surgical care measures were followed. Subsequently, the hospital reports implementing enhanced patient-focused infection prevention measures.