

Fax Page 1 To:

Jefferson County Public Health

Adapted from WA DOH Form 347-102 (360) 385-3878 (Confidential FAX line)

CONFIDENTIAL SEXUALLY TRANSMITTED INFECTION CASE REPORT Report STIs within 3 work days (WAC 246-101-101/301)

DATIENT INICO	DAAATION		,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			,. (4C 240 101 10	, , , ,				
PATIENT INFORMATION				FIRST NAME			MIDDLE NAME DATE OF BIRTH					
LAST NAME FII			FIRST NAME						DATE OF BIRTH			
455555 /	7					01777			MO		DAY	YR
ADDRESS (Unhoused or unstably housed in the pa				st 3 mont	:hs)	CITY				STATE	ZIP CO	DE
TELEPHONE		EMAIL		E	NGLIS	SH SPEAKING	G? ☐ Yes ☐ N	o *instructi		DIAGNOSI	S DATE	
()		1				red Languag		pg. 9	- 1	1 on	DAY	YR
SEX ASSIGNED	GENDER IDENT	ITY		ETHNICIT			RACE CATEGO	RY (check				Instructions on page 3
AT BIRTH	☐ Male	☐ Transgend	er MTF			Latina/o/x	☐ White	,		,	☐ Asia	an
☐ Male ☐ Female ☐ Transgende		· I — '			□ Black □ American Indian / Al □ Native Hawaiian / Of							
☐ Female ☐ Nonbinary / ☐ Other		Unknown										
☐ Intersex ☐ Refused	Genderque	er 🗌 Refused		☐ Refused								
CURRENTLY	DEACON FOR E	VAR4 (also also assa).	CENIE	SER OF CE	V DAD	ENIEDC (.ll						<u> </u>
PREGNANT?		XAM (check one):	1		X PAR	Transge	call that apply):				/AIDS Case Repor	CURRENTLY ON PrEP?
☐ Yes	☐ Exposed to I		│			☐ Transge	-		ious positive HIV diagnosis at this visit* ☐ Yes			
□ No		ຕ m (No Symptoms)	I —		_		l l			e HIV test at this visit		I .
□ Unk. □ NA		(nderquee	er	Unknov	vn	☐ Did r	ot test	(unknow	n status)	□NA
DIAGNOSIS - D	DISEASE											
GONORRHEA (I	ab confirmed)								SYPHILIS			
DIAGNOSIS (che	eck one)	SITES (all that	apply): ¦	TREATME	ENT (c	heck all pres	scribed):			check on		
Asymptomati	Cervix		☐ Ceftriaxone: ☐ 250 mg ☐ 500 mg ☐ 1 g				Primary (Chancre, etc.)					
☐ Symptomatic, Uncomplicated☐ Pelvic Inflammatory Disease		☐ Urethra☐ Urine		☐ Cefixime: ☐ 400 mg ☐ 800 mg ☐ Azithromycin: ☐ 1 g ☐ 2 g				☐ Secondary (Rash, etc.) ☐ Early Latent (< 1 year)				
☐ Ophthalmia ☐ Rectu						ne: \square 100 mg BID x 7 days			☐ Unknown Duration or Late			
Disseminated		☐ Pharynx	☐ Gentamicin:] [☐ Cong	enital		
Other Compli	caπons:	¦ ☐ Vagina	☐ Gemifloxacin: ☐ 320 mg			g	MANIFESTATIONS (check all that apply			that apply):		
Data Tastadi		¦ ☐ Ocular ! ☐ Other:	Othor		Other:			☐ Neurologic ☐ Otic ☐ Ocular ☐ Tertiary				
Date Tested:	h soufiumed)			Date Pres	scribed	u:				AENIT / - I-		
CHLAMYDIA (lab confirmed)		SITES (all that apply):		TREATMENT (check all prescribed):				TREATMENT (check one):				
DIAGNOSIS (check one) ☐ Asymptomatic ☐ Symptomatic, Uncomplicated		☐ Cervix		☐ Azithromycin: ☐ 1 g ☐ Doxycycline: ☐ 100 mg BID x 7 days				Bicillin L - A: ☐ 2.4 MU IM x 1 ☐ 2.4 MU IM x 3 Doxycycline: ☐ 100 mg BID x 14 days				
							- 1,					
Pelvic Inflami	matory Disease	☐ Urine		☐ Levofloxacin: ☐ 500 mg daily x 7 days				100 mg BID x 28 days				
☐ Ophthalmia☐ Other Complications:		Rectum		Other:			1	Benzathine				
		☐ Pharynx ☐ Vagina] 				PCN-G: 50,000 units/kg IM x 3				
		☐ Ocular							Aqueou	IS 🗌 1	18-24 MU/da	ay IV
Date Tested: Other:			Date Prescribed:				1.	Crystalline for 10-14 days Penicillin G:				
HERPES SIMPLE					OTHER	DISEASES						
DIAGNOSIS Genital (initia	l infection only)	LABORATORY CO	INFIRMA	TION	Cha	ncroid nuloma Ingui	nale		otner: _			
☐ Neonatal	inicotion omy,	☐ No					na Venereum		Date Pr	escribed:		
PARTNER TREA	ATMENT PLAN	(check one or r	nore op	tions)								
Providers should manage partner treatment by either treating partners in-person or by prescribing medication for patients to give to their sex partners (see side 2												
for additional info	•	of partners treats	nd followir	na medical	l ovalu	ation:	· · · · · · · · · · · · · · · · · · ·	Tu	rn over f	or Partner T	reatment Plan	Instructions
☐ In-person evaluation - Number of partners treated following medical evaluation:												
delivered by the patient to their partner(s): *Patient-delivered treatment is not recommended for men who have sex with men or patients with syphilis												
REPORTING CL												
DATE	FACILITY NA	ME				DIAGN	IOSING CLINICIA	AN				
ADDRESS	·					CITY			STATE		ZIP	
PERSON COMPLETING FORM TELEPHONE EMAIL												
	-			())							

PARTNER MANAGEMENT PLAN INSTRUCTIONS

Gonorrhea or Chlamydia Infection: Partner Treatment

All partners should be treated as if they are infected.

The Washington State Department of Health strongly encourages providers to take responsibility to ensure partner treatment for heterosexuals, by examining and treating all patient's sex partners from the previous 60 days.

If an examination is not possible, providers should offer medication for all sex partners whom patients are able to contact.

Jefferson County Public Health may be able to provide free medication to your patient to give their partner(s), if resources permit. Please contact your local health jurisdiction to report cases and inquire about partner management resources, possibly including EPT medications.

Jefferson County Public Health recommends you refer all **MSM patients** and **all patients with syphilis or newly diagnosed HIV** to the health department for help notifying partners to ensure that partners receive medication; the opportunity to test for HIV, syphilis, gonorrhea, and chlamydia; and evaluation for HIV Pre-Exposure Prophylaxis (PrEP). Please inform the patient that the health department will contact them to assist with partner notification.

Although the Health Department requests that you refer patients with these risks to us, we also ask that you make every effort to help patients assure that their partners are treated.

Complete the partner management plan on the Confidential Sexually Transmitted Infection Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call the Jefferson County Public Health: (360) 385-9400.

Other STIs: Partner Treatment

All patients with infectious syphilis, chancroid, LGV, or granuloma inguinale are routinely contacted by public health staff. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing. Per CDC treatment guidelines, sex partners of patients who are diagnosed with early syphilis (primary, secondary, or early latent) and may be incubating disease should be treated regardless of test results. Alternative treatment for penicillin allergy among non-pregnant patients, such as an appropriate dosage of doxycycline, can also be found in the guidelines.

RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON REPORTS*

GONORRHEA -- Uncomplicated

Ceftriaxone 500 mg IM as a single dose for persons weighing < 150 kg (330 lbs), or 1 g IM as a single dose for persons weighing > 150 kg (330 lbs) †

Alternatives for uncomplicated infections of the cervix, urethra, or rectum: ‡

Gentamicin 240 mg IM as a single dose PLUS azithromycin 2 g PO as a single dose $\bf OR$ Cefixime 800 mg orally as a single dose †

- † If treating with ceftriaxone or cefixime, and chlamydia infection has not been excluded, providers should treat for chlamydia with doxycycline 100 mg PO BID for 7 days. During pregnancy, azithromycin 1 g PO as a single dose is recommended to treat chlamydia.
- [‡] Ceftriaxone is the only recommended treatment for pharyngeal gonorrhea. Consult an infectious disease provider or other STI expert for assistance if alternative treatment is required.

CHLAMYDIA -- Uncomplicated

Doxycycline 100 mg PO BID for 7 days **OR** Azithromycin 1g PO as a single dose

Alternatives:

Erythromycin (base) 500 mg PO QID for 7 days OR

Ethylsuccinate 800 mg PO QID for 7 days OR

Ofloxacin 300 mg PO BID for 7 days OR

Levofloxacin 500 mg PO for 7 days

SYPHILIS -- PRIMARY, SECONDARY, OR EARLY LATENT (<1 YEAR)

Benzathine penicillin G 2.4 million units IM in a single dose

SYPHILIS -- LATE OR UNKNOWN DURATION

Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals

DOH 347-102, updated 12/14/2022. For persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).

^{*} Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (https://www.cdc.gov/std/treatment/default.htm) for further information on treating pregnant patients, infections of the pharynx, patients with allergies, treatment of infants, and other details.

List of Preferred Languages:

Instructions: Complete the Preferred Language section if the patient's preferred language is not English. Patient's preferred language shall be identified by the patient and reported using one of the below categories.

Language Identity Codes:

(L18) Karen	(L35) Sign languages
(L19) Khmer/Cambodian	(L36) Somali
(L20) Kinyarwanda	(L37) Spanish/Castilian
(L21) Korean	(L38) Swahili/Kiswahili
(L22) Kosraean	(L39) Tagalog
(L23) Lao	(L40) Tamil
(L24) Mandarin	(L41) Telugu
(L25) Marshallese	(L42) Thai
(L26) Mixteco	(L43) Tigrinya
(L27) Nepali	(L44) Ukrainian
(L28) Oromo	(L45) Urdu
(L29) Panjabi/Punjabi	(L46) Vietnamese
(L30) Pashto	(L77) Other language
(L31) Portuguese	(L88) Patient declined to respond
(L32) Romanian/Rumanian	(L99) Unknown
	(L19) Khmer/Cambodian (L20) Kinyarwanda (L21) Korean (L22) Kosraean (L23) Lao (L24) Mandarin (L25) Marshallese (L26) Mixteco (L27) Nepali (L28) Oromo (L29) Panjabi/Punjabi (L30) Pashto (L31) Portuguese

(L33) Russian

(L34) Samoan

Race Category and Identity Instructions:

Instructions: The patient's race may either be reported in the general Race Category section or the Extended Race section on page 1 of the case report. When completing the Extended Race section, please report using the codes provided in the list below. If the patient self-identifies as more than one race, each race code shall be reported within the extended race section. Unknown race can be documented within either one of the race sections.

Extended Race Codes:

(L16) Hmong

(L17) Japanese

(R01) Afghan	(R22) Fijian	(R43) Mexican/Mexican American	(R64) Tongan
(R02) Afro-Caribbean	(R23) Filipino	(R44) Middle Eastern	(R65) Ugandan
(R03) Alaska Native	(R24) First Nations	(R45) Mien	(R66) Ukrainian
(R04) American Indian	(R25) Guamanian or Chamorro	(R46) Moroccan	(R67) Vietnamese
(R05) Arab	(R26) Hmong/Mong	(R47) Native Hawaiian	(R68) White
(R06) Asian	(R27) Indigenous-Latina/o/x	(R48) Nepalese	(R69) Yemeni
(R07) Asian Indian	(R28) Indonesian	(R49) North African	(R77) Other race
(R08) Bamar/Burman/Burmese	(R29) Iranian	(R50) Oromo	(R99) Unknown
(R09) Bangladeshi	(R30) Iraqi	(R51) Pacific Islander	
(R10) Bhutanese	(R31) Japanese	(R52) Pakistani	
(R11) Black or African American	(R32) Jordanian	(R53) Puerto Rican	
(R12) Central American	(R33) Karen	(R54) Romanian/Rumanian	
(R13) Cham	(R34) Kenyan	(R55) Russian	
(R14) Chicana/o or Chicanx	(R35) Khmer/Cambodian	(R56) Samoan	
(R15) Chinese	(R36) Korean	(R57) Saudi Arabian	
(R16) Congolese	(R37) Kuwaiti	(R58) Somali	
(R17) Cuban	(R38) Lao	(R59) South African	
(R18) Dominican	(R39) Lebanese	(R60) South American	
(R19) Egyptian	(R40) Malaysian	(R61) Syrian	
(R20) Eritrean	(R41) Marshallese	(R62) Taiwanese	
(R21) Ethiopian	(R42) Mestizo	(R63) Thai	