

Fax Page 1 To:

Skagit County Department of Health (360) 416-1515 (Confidential FAX line)

Adapted from WA DOH Form 347-102

CONFIDENTIAL SEXUALLY TRANSMITTED INFECTION CASE REPORT
Report STIs within 3 work days (WAC 246-101-101/301)

PATIENT INFO	RMATION	·										
LAST NAME		FIRST NAME					MIDDLE NAME			DATE OF BIRTH		
									мо	D	AY	YR
ADDRESS (Unhoused or unstably housed in the pa				months)	CITY					STATE	ZIP COD	E
TELEPHONE		EMAIL		ENGL	ISH SP	EAKING	G? 🗌 Yes 🔲 N	o *instruct		IAGNOSIS	DATE	
()				Prefe	rred La	inguag	e (Code*: L			10 1	DAY	YR
SEX ASSIGNED	GENDER IDENT	TITY	ET	HNICITY			RACE CATEGO	RY (check	all that a	apply)*:		structions on page 3
AT BIRTH ☐ Male ☐ Transgender						☐ White ☐ Asian ☐ Other						
☐ Male ☐ Female ☐ Transgender F		I				☐ American Ir	ndian / Ala	iska Nati	ive	Unkn		
☐ Female ☐ Nonbinary / ☐ Other ☐ Intersex ☐ Genderqueer ☐ Refused			Refused			☐ Native Hawaiian / Other Pacific Islander ☐ Refused						
☐ Refused			_	EXTENDED RACE COL			CE CODE	E(S)*: R R R R				
CURRENTLY	REASON FOR E	XAM (check one):	GENDER	OF SEX PA	RTNERS	S (checl	all that apply):	HIV STA	rus	*Submit HIV/A	IDS Case Report	CURRENTLY
PREGNANT? Exposed to I				_ 0			· ·			ON PrEP?		
☐ Yes ☐ No	Symptomatic		☐ Femal	_ 0		• • • • • • • • • • • • • • • • • • •			HIV diagnosis at this visit* ☐ Yes			
Unk. 🗆 NA	☐ Routine Exar	n (No Symptoms)	S) Nonbinary / Genderqueer			_ -			ative HIV test at this visit			
DIAGNOSIS - D	DISFASE			11							,	
GONORRHEA (la									SYPHILI	s		
DIAGNOSIS (che	eck one)	SITES (all that a	pply): TR	TREATMENT (check all prescribed):				STAGE (check one):				
Asymptomati		☐ Cervix					g 🗌 500 mg		☐ Primary (Chancre, etc.)			
Symptomatic, Uncomplicated		1		Cefixime:			g 🔲 800 mg		Secondary (Rash, etc.)			
☐ Pelvic Inflammatory Disease☐ Ophthalmia				☐ Azithromycin: ☐ 1 g ☐ 2 g				☐ Early Latent (< 1 year) ☐ Unknown Duration or Late				
☐ Disseminated		Pharynx		☐ Doxycycline: ☐ 100 mg BID x 7 days ☐ Gentamicin: ☐ 240 mg					☐ Congenital			
☐ Other Compli	cations:	· ' - ' - ' - ' - ' - ' - ' - ' - ' - '		☐ Gemifloxacin: ☐ 320 mg					MANIFESTATIONS (check all that apply):			
		☐ Ocular ☐		Other:				Neurologic ☐ Otic ☐ Ocular ☐ Tertiary				
Date Tested:		Other:	—— Da	te Prescrib	ed:				Treationogie Cottle Cottain Circlinary			
CHLAMYDIA (la	· · · · · · · · · · · · · · · · · · ·								TREATMENT (check one):			
DIAGNOSIS (che	-	SITES (all that apply):						Bicillin L - A: ☐ 2.4 MU IM x 1				
☐ Asymptomati ☐ Symptomatic				☐ Azithromycin: ☐ 1 g ☐ Doxycycline: ☐ 100 mg BID x 7 days				☐ 2.4 MU IM x 3				
	matory Disease			Levofloxacin: 500 mg daily x 7 days				Doxycycline: ☐ 100 mg BID x 14 days ☐ 100 mg BID x 28 days				
☐ Ophthalmia		☐ Rectum		☐ Other:				Benzathine 50,000 units/kg IM x 1				
☐ Other Complications:		☐ Pharynx ☐ Vagina						PCN-G: 50,000 units/kg IM x 3				
		☐ Vagilla ☐ Ocular							Aqueou	S □ 18	3-24 MU/day	IV
				Date Prescribed:				Crystalline for 10-14 days				
HERPES SIMPLE	X			OTHE	R DISE	ASES			Penicilli	n G:		
DIAGNOSIS		LABORATORY CO	NFIRMATIO		ancroid				Other: _			
☐ Genital (initia☐ Neonatal	I infection only)	☐ Yes ☐ No			anulom		nale na Venereum		Date Pro	escribed:		
	ATMENT PLAN	(check one or m	ore option	<u> </u>	ПРПОВІ	41141611	Ta veneream					
		treatment by either			rson or	by pre	scribing medicat	ion for pa	tients to	give to the	ir sex partne	rs (see side 2
for additional information).												
In-person evaluation - Number of partners treated following medical evaluation:												
☐ Patient-delivered treatment* - Number of partners for whom provider prescribed or provided expedited partner therapy (EPT) medication pack to be delivered by the patient to their partner(s): *Patient-delivered treatment is not recommended for men who have sex with men or patients with syphilis												
REPORTING CLINIC INFORMATION												
DATE	FACILITY NA	AME				DIAGN	IOSING CLINICIA	AN				
ADDRESS					CITY				STATE		ZIP	
PERSON COMPLETING FORM				TELEPHO	NE N			EMA	IL			

PARTNER MANAGEMENT PLAN INSTRUCTIONS

Gonorrhea or Chlamydia Infection: Partner Treatment

All partners should be treated as if they are infected.

The Washington State Department of Health strongly encourages providers to take responsibility to ensure partner treatment for heterosexuals, by examining and treating all patient's sex partners from the previous 60 days.

If an examination is not possible, providers should offer medication for all sex partners whom patients are able to contact.

Skagit County Department of Health may be able to provide free medication to your patient to give their partner(s), if resources permit. Please contact your local health jurisdiction to report cases and inquire about partner management resources, possibly including EPT medications.

Skagit County Department of Health recommends you refer all MSM patients and all patients with syphilis or newly diagnosed HIV to the health department for help notifying partners to ensure that partners receive medication; the opportunity to test for HIV, syphilis, gonorrhea, and chlamydia; and evaluation for HIV Pre-Exposure Prophylaxis (PrEP). Please inform the patient that the health department will contact them to assist with partner notification.

Although the Health Department requests that you refer patients with these risks to us, we also ask that you make every effort to help patients assure that their partners are treated.

Complete the partner management plan on the Confidential Sexually Transmitted Infection Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call the Skagit County Department of Health: (360) 416-1500.

Other STIs: Partner Treatment

All patients with infectious syphilis, chancroid, LGV, or granuloma inguinale are routinely contacted by public health staff. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing. Per CDC treatment guidelines, sex partners of patients who are diagnosed with early syphilis (primary, secondary, or early latent) and may be incubating disease should be treated regardless of test results. Alternative treatment for penicillin allergy among non-pregnant patients, such as an appropriate dosage of doxycycline, can also be found in the guidelines.

RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON REPORTS*

GONORRHEA -- Uncomplicated

Ceftriaxone 500 mg IM as a single dose for persons weighing < 150 kg (330 lbs), or 1 g IM as a single dose for persons weighing > 150 kg (330 lbs) †

Alternatives for uncomplicated infections of the cervix, urethra, or rectum: ‡

Gentamicin 240 mg IM as a single dose PLUS azithromycin 2 g PO as a single dose $\bf OR$ Cefixime 800 mg orally as a single $\bf dose^{\dagger}$

- † If treating with ceftriaxone or cefixime, and chlamydia infection has not been excluded, providers should treat for chlamydia with doxycycline 100 mg PO BID for 7 days. During pregnancy, azithromycin 1 g PO as a single dose is recommended to treat chlamydia.
- [‡] Ceftriaxone is the only recommended treatment for pharyngeal gonorrhea. Consult an infectious disease provider or other STI expert for assistance if alternative treatment is required.

CHLAMYDIA -- Uncomplicated

Doxycycline 100 mg PO BID for 7 days **OR**Azithromycin 1g PO as a single dose

Alternatives:

Erythromycin (base) 500 mg PO QID for 7 days ${\bf OR}$

Ethylsuccinate 800 mg PO QID for 7 days OR

Ofloxacin 300 mg PO BID for 7 days OR

Levofloxacin 500 mg PO for 7 days

SYPHILIS -- PRIMARY, SECONDARY, OR EARLY LATENT (<1 YEAR)

Benzathine penicillin G 2.4 million units IM in a single dose

SYPHILIS -- LATE OR UNKNOWN DURATION

Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals

DOH 347-102, updated 12/14/2022. For persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).

^{*} Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (https://www.cdc.gov/std/treatment/default.htm) for further information on treating pregnant patients, infections of the pharynx, patients with allergies, treatment of infants, and other details.

List of Preferred Languages:

Instructions: Complete the Preferred Language section if the patient's preferred language is not English. Patient's preferred language shall be identified by the patient and reported using one of the below categories.

Language Identity Codes:

(L01) Amharic	(L18) Karen	(L35) Sign languages
(L02) Arabic	(L19) Khmer/Cambodian	(L36) Somali
(LO3) Balochi/Baluchi	(L20) Kinyarwanda	(L37) Spanish/Castilian
(L04) Burmese	(L21) Korean	(L38) Swahili/Kiswahili
(L05) Cantonese	(L22) Kosraean	(L39) Tagalog
(L06) Chinese (unspecified)	(L23) Lao	(L40) Tamil
(L07) Chamorro	(L24) Mandarin	(L41) Telugu
(L08) Chuukese	(L25) Marshallese	(L42) Thai
(L09) Dari	(L26) Mixteco	(L43) Tigrinya
(L10) Farsi/Persian	(L27) Nepali	(L44) Ukrainian
(L11) Fijian	(L28) Oromo	(L45) Urdu
(L12) Filipino/Pilipino	(L29) Panjabi/Punjabi	(L46) Vietnamese
(L13) French	(L30) Pashto	(L77) Other language
(L14) German	(L31) Portuguese	(L88) Patient declined to respond
(L15) Hindi	(L32) Romanian/Rumanian	(L99) Unknown

(L33) Russian

(L34) Samoan

Race Category and Identity Instructions:

Instructions: The patient's race may either be reported in the general Race Category section or the Extended Race section on page 1 of the case report. When completing the Extended Race section, please report using the codes provided in the list below. If the patient self-identifies as more than one race, each race code shall be reported within the extended race section. Unknown race can be documented within either one of the race sections.

Extended Race Codes:

(L16) Hmong

(L17) Japanese

(R01) Afghan	(R22) Fijian	(R43) Mexican/Mexican American	(R64) Tongan
(R02) Afro-Caribbean	(R23) Filipino	(R44) Middle Eastern	(R65) Ugandan
(R03) Alaska Native	(R24) First Nations	(R45) Mien	(R66) Ukrainian
(R04) American Indian	(R25) Guamanian or Chamorro	(R46) Moroccan	(R67) Vietnamese
(R05) Arab	(R26) Hmong/Mong	(R47) Native Hawaiian	(R68) White
(R06) Asian	(R27) Indigenous-Latina/o/x	(R48) Nepalese	(R69) Yemeni
(R07) Asian Indian	(R28) Indonesian	(R49) North African	(R77) Other race
(R08) Bamar/Burman/Burmese	(R29) Iranian	(R50) Oromo	(R99) Unknown
(R09) Bangladeshi	(R30) Iraqi	(R51) Pacific Islander	
(R10) Bhutanese	(R31) Japanese	(R52) Pakistani	
(R11) Black or African American	(R32) Jordanian	(R53) Puerto Rican	
(R12) Central American	(R33) Karen	(R54) Romanian/Rumanian	
(R13) Cham	(R34) Kenyan	(R55) Russian	
(R14) Chicana/o or Chicanx	(R35) Khmer/Cambodian	(R56) Samoan	
(R15) Chinese	(R36) Korean	(R57) Saudi Arabian	
(R16) Congolese	(R37) Kuwaiti	(R58) Somali	
(R17) Cuban	(R38) Lao	(R59) South African	
(R18) Dominican	(R39) Lebanese	(R60) South American	
(R19) Egyptian	(R40) Malaysian	(R61) Syrian	
(R20) Eritrean	(R41) Marshallese	(R62) Taiwanese	
(R21) Ethiopian	(R42) Mestizo	(R63) Thai	