

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2013^{1, 2, 3}
October 8, 2014 download of January 2013 – December 2013 NHSN data

| Hospitals | Type of Surgical Procedure ⁴ | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|---|---|---------------|----------------------|------------------|---------------|----------------------|--|---------------|----------------------|------------------------|---------------|----------------------|------------------|---------------|----------------------|------------------------------|---------------|----------------------|--------------|---------------|----------------------|-------------------------------|---------------|----------------------|--------------|---------------|----------------------|--------------|---------------|-------|------|-------|-------|------|
| | Abdominal Hysterectomy | | | Colon Surgery | | | Coronary Artery Bypass Graft ⁵ | | | | | | Heart Surgery | | | Hip Replacement ⁵ | | | | | | Knee Replacement ⁵ | | | | | | | | | | | | |
| | | | | | | | Chest and Secondary Incision Site | | | Chest Incision Only | | | | | | Overall | | | Initial | | Revision | | Overall | | Initial | | Revision | | Overall | | | | | |
| | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | | | | | |
| Capital Medical Center | 0.00 | 3.45 | 29 | 0.00 | 0.00 | 64 | | | | | | | | | | | | | 0.00 | 0.55 | 182 | 0.00 | 0.00 | 25 | 0.00 | 0.48 | 0.00 | 0.00 | 169 | 4.17 | 4.17 | 24 | 0.52 | 0.52 |
| Cascade Valley Hospital | 0.00 | 0.00 | 2 | 2.63 | 2.63 | 38 | | | | | | | | | | | | | 0.00 | 0.00 | 26 | - | - | 0 | 0.00 | 0.00 | 0.00 | 0.00 | 41 | 0.00 | 0.00 | 1 | 0.00 | 0.00 |
| Central Washington Hospital ¹⁴ | 0.00 | 0.00 | 3 | 5.68* | 10.23* | 176* | 0.86 | 4.31 | 116 | 0.00 | 0.00 | 8 | 0.81 | 4.03 | 0.00 | 1.10 | 91 | 1.23 | 1.23 | 244 | 0.00 | 0.00 | 22 | 1.13 | 1.13 | 0.00 | 0.49 | 406 | 3.85 | 3.85 | 26 | 0.23 | 0.69 | |
| Coulee Medical Center | 0.00 | 0.00 | 11 | 0.00 | 0.00 | 2 | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Deaconess Medical Center ¹⁸ | 0.99 | 0.99 | 101 | 5.04 | 8.63 | 139 | 3.49 | 6.98 | 86 | 0.00 | 0.00 | 25 | 2.70 | 5.41 | 0.00 | 0.00 | 52 | 0.49 | 1.47 | 204 | 5.00 | 5.00 | 20 | 0.89 | 1.79 | 0.52 | 0.52 | 191 | 4.00 | 4.00 | 25 | 0.93 | 0.93 | |
| Evergreen Healthcare | 0.00 | 1.24 | 161 | 4.84 | 6.45 | 62 | | | | | | | | | | | | 0.44 | 0.88 | 228 | 0.00 | 0.00 | 14 | 0.41 | 0.83 | 0.00 | 0.00 | 450 | 2.70 | 2.70 | 37 | 0.21 | 0.21 | |
| Forks Community Hospital | - | - | 0 | | | | | | | | | | | | | | | 0.00 | 0.00 | 2 | - | - | 0 | 0.00 | 0.00 | 0.00 | 0.00 | 6 | - | - | 0 | 0.00 | 0.00 | |
| Grays Harbor Community Hospital ⁶ | 4.76 | 4.76 | 21 | 0.00 | 0.00 | 24 | | | | | | | | | | | | 0.00 | 2.38 | 42 | 0.00 | 0.00 | 2 | 0.00 | 2.27 | 0.00 | 1.52 | 66 | 0.00 | 0.00 | 1 | 0.00 | 1.49 | |
| Harborview Medical Center ^{7, 25} | 0.00 | 5.56 | 36 | 3.23 | 6.45 | 62 | - | - | 0 | - | - | 0 | - | - | 0.00 | 0.00 | 11 | 0.00 | 2.94 | 34 | - | - | 0 | 0.00 | 2.94 | 0.00 | 0.00 | 1 | 25.00 | 25.00 | 4 | 20.00 | 20.00 | |
| Harrison Medical Center | 0.00 | 0.00 | 71 | 3.11 | 6.67 | 225 | 1.08 | 2.70 | 185 | 0.00 | 0.00 | 1 | 1.08 | 2.69 | 0.00 | 0.00 | 118 | 0.42 | 0.42 | 237 | 0.00 | 0.00 | 11 | 0.40 | 0.40 | 0.28 | 0.28 | 358 | 0.00 | 0.00 | 14 | 0.27 | 0.27 | |
| Highline Medical Center | 0.00 | 0.00 | 43 | 4.05 | 5.41 | 74 | | | | | | | | | | | | 0.00 | 0.00 | 95 | 0.00 | 0.00 | 6 | 0.00 | 0.00 | 0.00 | 0.00 | 124 | 0.00 | 0.00 | 5 | 0.00 | 0.00 | |
| Island Hospital | 0.00 | 0.00 | 21 | 0.00 | 3.03 | 33 | | | | | | | | | | | | 0.00 | 0.00 | 94 | 0.00 | 0.00 | 3 | 0.00 | 0.00 | 0.56 | 0.56 | 179 | 0.00 | 0.00 | 12 | 0.52 | 0.52 | |
| Jefferson General Hospital ^{22, 26} | 0.00 | 0.00 | 3 | 0.00 | 9.09 | 11 | | | | | | | | | | | | 6.45 | 12.90 | 31 | 0.00 | 0.00 | 1 | 6.25 | 12.50 | 0.00 | 4.35 | 23 | 0.00 | 0.00 | 2 | 0.00 | 4.00 | |
| Kadlec Medical Center | 1.09 | 2.17 | 92 | 3.33 | 4.67 | 150 | 1.53 | 1.53 | 261 | 11.11 | 11.11 | 9 | 1.85 | 1.85 | 0.00 | 0.00 | 100 | 0.00 | 0.00 | 142 | 0.00 | 9.09 | 11 | 0.00 | 0.65 | 0.00 | 0.00 | 147 | 0.00 | 0.00 | 13 | 0.00 | 0.00 | |
| Kennewick General Hospital ^{8, 27} | 4.00 | 8.00 | 50* | 2.86 | 8.57 | 35 | | | | | | | | | | | | 0.00 | 0.00 | 71 | 0.00 | 0.00 | 5 | 0.00 | 0.00 | 1.53 | 1.53 | 131 | 0.00 | 0.00 | 12 | 1.40 | 1.40 | |

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2013^{1, 2, 3}
October 8, 2014 download of January 2013 – December 2013 NHSN data

| Hospitals | Type of Surgical Procedure ⁴ | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|---------------------------------------|---|------------|-------------------|------------------|------------|-------------------|--|------------|-------------------|------------------------|------------|-------------------|------------------|------------|-------------------|------------------------------|------------|-------------------|-----------|------------|-------------------|-------------------------------|------------|-------------------|-----------|------------|-------------------|-----------|------------|------|---------|------|------|
| | Abdominal Hysterectomy | | | Colon Surgery | | | Coronary Artery Bypass Graft ⁵ | | | | | | Heart Surgery | | | Hip Replacement ⁵ | | | | | | Knee Replacement ⁵ | | | | | | | | | | | |
| | | | | | | | Chest and Secondary Incision Site | | | Chest Incision Only | | | | | | Overall | Initial | | | Revision | | | Overall | | Initial | | | Revision | | | Overall | | |
| | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | | | | |
| Kittitas Valley Community Hospital | 0.00 | 0.00 | 2 | 0.00 | 0.00 | 16 | | | | | | | | | | | | 0.00 | 1.56 | 64 | 0.00 | 0.00 | 2 | 0.00 | 1.52 | 0.00 | 1.23 | 81 | 0.00 | 0.00 | 3 | 0.00 | 1.19 |
| Lake Chelan Hospital | - | - | 0 | 0.00 | 0.00 | 3 | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Legacy Salmon Creek Hospital | 1.92 | 1.92 | 104 | 2.16 | 2.88 | 139 | | | | | | | | | | | | 0.49 | 0.49 | 206 | 0.00 | 0.00 | 10 | 0.46 | 0.46 | 0.00 | 0.00 | 610 | 0.00 | 0.00 | 34 | 0.00 | 0.00 |
| Lincoln Hospital | 0.00 | 0.00 | 5 | 0.00 | 0.00 | 10 | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Lourdes Health Network | 0.00 | 0.00 | 43 | 0.00 | 0.00 | 8 | | | | | | | | | | | | 0.00 | 0.00 | 105 | 0.00 | 0.00 | 25 | 0.00 | 0.00 | 0.77 | 0.77 | 390 | 0.00 | 0.00 | 36 | 0.70 | 0.70 |
| Mary Bridge Children's Hospital | | | | 0.00 | 3.70 | 27 | | | | | | | | 0.00 | 0.00 | 10 | - | - | 0 | - | - | 0 | - | - | - | - | 0 | - | - | 0 | - | - | |
| Mason General Hospital | 0.00 | 0.00 | 8 | 9.09 | 9.09 | 11 | | | | | | | | | | | | 0.00 | 0.00 | 33 | 0.00 | 0.00 | 1 | 0.00 | 0.00 | 0.00 | 0.00 | 41 | - | - | 0 | 0.00 | 0.00 |
| Mid-Valley Hospital | 0.00 | 0.00 | 6 | 0.00 | 0.00 | 6 | | | | | | | | | | | | 0.00 | 0.00 | 18 | - | - | 0 | 0.00 | 0.00 | 0.00 | 0.00 | 23 | - | - | 0 | 0.00 | 0.00 |
| Morton General Hospital | - | - | 0 | 0.00 | 0.00 | 2 | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Multicare Auburn Medical Center | 0.00 | 0.00 | 52 | 0.00 | 0.00 | 45 | | | | | | | | | | | | 0.00 | 0.00 | 84 | 0.00 | 0.00 | 4 | 0.00 | 0.00 | 0.00 | 0.85 | 118 | 0.00 | 0.00 | 10 | 0.00 | 0.78 |
| Multicare Good Samaritan Hospital | 0.74 | 0.74 | 269 | 2.53 | 3.16 | 158 | | | | | | | | | | | | 0.42 | 0.42 | 238 | 0.00 | 0.00 | 16 | 0.39 | 0.39 | 0.00 | 0.39 | 254 | 0.00 | 0.00 | 10 | 0.00 | 0.38 |
| Newport Community Hospital | | | | 0.00 | 0.00 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| North Valley Hospital | - | - | 0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Northwest Hospital ¹⁹ | 0.89 | 1.34 | 224 | 2.68 | 3.57 | 112 | 3.03 | 3.03 | 33 | 0.00 | 0.00 | 3 | 2.78 | 2.78 | 0.00 | 0.00 | 38 | 0.00 | 0.71 | 281 | 2.44 | 2.44 | 41 | 0.31 | 0.93 | 0.77 | 1.03 | 390 | 0.00 | 0.00 | 34 | 0.71 | 0.94 |
| Ocean Beach Hospital | | | | 0.00 | 0.00 | 3 | | | | | | | | | | | | - | - | 0 | 0.00 | 0.00 | 1 | 0.00 | 0.00 | 0.00 | 0.00 | 4 | - | - | 0 | 0.00 | 0.00 |

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2013^{1, 2, 3}
October 8, 2014 download of January 2013 – December 2013 NHSN data

| Hospitals | Type of Surgical Procedure ⁴ | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|--|---|---------------|----------------------|------------------|---------------|----------------------|--|---------------|----------------------|------------------------|---------------|----------------------|------------------|---------------|----------------------|------------------------------|---------------|----------------------|--------------|---------------|----------------------|-------------------------------|---------------|----------------------|--------------|---------------|----------------------|--------------|---------------|----------------------|------|------|------|
| | Abdominal Hysterectomy | | | Colon Surgery | | | Coronary Artery Bypass Graft ⁵ | | | | | | Heart Surgery | | | Hip Replacement ⁵ | | | | | | Knee Replacement ⁵ | | | | | | | | | | | |
| | | | | | | | Chest and Secondary Incision Site | | | Chest Incision Only | | | | | | Overall | Initial | | | Revision | | | Overall | Initial | | | Revision | | | Overall | | | |
| | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | | | |
| Olympic Medical Center | 0.00 | 0.00 | 28 | 5.66 | 7.55 | 53 | | | | | | | | | | | 0.81 | 0.81 | 124 | 0.00 | 0.00 | 3 | 0.79 | 0.79 | 0.00 | 0.00 | 155 | - | - | 0 | 0.00 | 0.00 | |
| Othello Community Hospital | 0.00 | 0.00 | 2 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Overlake Hospital Medical Center | 1.24 | 1.98 | 404 | 2.19 | 4.74 | 274 | 0.00 | 1.01 | 99 | 0.00 | 0.00 | 6 | 0.00 | 0.95 | 0.00 | 0.00 | 72 | 0.18 | 0.36 | 550 | 1.28 | 1.28 | 78 | 0.32 | 0.48 | 0.15 | 0.15 | 648 | 2.56 | 2.56 | 39 | 0.29 | 0.29 |
| PeaceHealth Saint John Medical Center | 2.00 | 2.00 | 50 | 1.33 | 12.00 | 75 | | | | | | | | | | | 1.04 | 1.55 | 193 | 0.00 | 7.14 | 14 | 0.97 | 1.93 | 0.00 | 0.00 | 273 | 0.00 | 0.00 | 14 | 0.00 | 0.00 | |
| PeaceHealth Saint Joseph Medical Center ²⁰ | 0.00 | 0.00 | 93 | 6.17 | 8.64 | 162 | 2.11 | 3.16 | 95 | 0.00 | 0.00 | 7 | 1.96 | 2.94 | 1.46 | 1.46 | 137 | 1.03 | 2.05 | 390 | 0.00 | 0.00 | 23 | 0.97 | 1.94 | 0.00 | 0.00 | 421 | 4.76 | 4.76 | 21 | 0.23 | 0.23 |
| PeaceHealth Southwest Medical Center | 0.35 | 0.35 | 288 | 1.98 | 3.47 | 202 | 0.46 | 0.92 | 217 | 0.00 | 0.00 | 1 | 0.46 | 0.92 | 0.00 | 0.00 | 124 | 0.53 | 0.53 | 379 | 0.00 | 0.00 | 53 | 0.46 | 0.46 | 0.65 | 0.65 | 306 | 0.00 | 0.00 | 22 | 0.61 | 0.61 |
| Prosser Memorial Hospital | 0.00 | 0.00 | 11 | 0.00 | 0.00 | 1 | | | | | | | | | | | 0.00 | 0.00 | 5 | - | - | 0 | 0.00 | 0.00 | 0.00 | 0.00 | 10 | 0.00 | 0.00 | 1 | 0.00 | 0.00 | |
| Providence Centralia Hospital | 2.08 | 2.08 | 48 | 4.29 | 5.71 | 70 | | | | | | | | | | | 0.73 | 0.73 | 137 | 0.00 | 0.00 | 10 | 0.68 | 0.68 | 0.60 | 0.60 | 166 | 0.00 | 0.00 | 9 | 0.57 | 0.57 | |
| Providence Everett Medical Center | 0.45 | 0.68 | 440 | 1.82 | 2.27 | 220 | 1.27 | 1.27 | 315 | 0.00 | 0.00 | 16 | 1.21 | 1.21 | 0.53 | 0.53 | 190 | 1.37 | 1.37 | 510 | 4.26 | 4.26 | 47 | 1.62 | 1.62 | 0.16 | 0.16 | 629 | 0.00 | 0.00 | 34 | 0.15 | 0.15 |
| Providence Holy Family Hospital ⁹ | 5.97 | 7.46 | 67* | 5.10 | 9.18 | 98 | | | | | | | | | | | 1.27 | 1.69 | 236 | 0.00 | 0.00 | 48 | 1.06 | 1.41 | 0.30 | 0.60 | 335 | 0.00 | 0.00 | 48 | 0.26 | 0.52 | |
| Providence Mount Carmel Hospital | 0.00 | 0.00 | 9 | 0.00 | 0.00 | 17 | | | | | | | | | | | 0.00 | 0.00 | 2 | - | - | 0 | 0.00 | 0.00 | 0.00 | 0.00 | 2 | 0.00 | 0.00 | 1 | 0.00 | 0.00 | |
| Providence Sacred Heart Medical Center | 0.53 | 0.53 | 376 | 3.68 | 4.82 | 353 | 0.51 | 0.51 | 396 | 0.00 | 0.00 | 18 | 0.48 | 0.48 | 0.00 | 0.00 | 533 | 0.32 | 0.96 | 628 | 1.03 | 1.03 | 97 | 0.41 | 0.97 | 0.14 | 0.14 | 731 | 0.00 | 0.00 | 99 | 0.12 | 0.12 |
| Providence Saint Mary Medical Center | 0.00 | 0.00 | 19 | 2.17 | 2.17 | 46 | | | | | | | | | | | 0.74 | 0.74 | 135 | 0.00 | 8.33 | 12 | 0.68 | 1.36 | 0.62 | 0.62 | 162 | 0.00 | 0.00 | 24 | 0.54 | 0.54 | |
| Providence Saint Peter Hospital | 0.00 | 0.00 | 101 | 0.97 | 1.45 | 207 | 0.00 | 1.33 | 226 | 0.00 | 0.00 | 14 | 0.00 | 1.25 | 0.00 | 0.65 | 155 | 0.19 | 0.39 | 513 | 1.35 | 2.70 | 74 | 0.34 | 0.68 | 0.12 | 0.36 | 826 | 0.80 | 0.80 | 125 | 0.21 | 0.42 |
| Pullman Regional Hospital ¹⁵ | 0.98 | 1.96 | 102 | 10.53 | 15.79 | 19 | | | | | | | | | | | 0.00 | 0.00 | 29 | 0.00 | 0.00 | 1 | 0.00 | 0.00 | 0.00 | 0.00 | 87 | 0.00 | 0.00 | 3 | 0.00 | 0.00 | |

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2013 ^{1, 2, 3}
October 8, 2014 download of January 2013 – December 2013 NHSN data

| Hospitals | Type of Surgical Procedure ⁴ | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|---|---|------------|-------------------|------------------|------------|-------------------|--|------------|-------------------|------------------------|------------|-------------------|------------------|------------|-------------------|------------------------------|------------|-------------------|-----------|------------|-------------------|-------------------------------|------------|-------------------|-----------|------------|-------------------|-----------|------------|---------|------|------|------|
| | Abdominal Hysterectomy | | | Colon Surgery | | | Coronary Artery Bypass Graft ⁵ | | | | | | Heart Surgery | | | Hip Replacement ⁵ | | | | | | Knee Replacement ⁵ | | | | | | | | | | | |
| | | | | | | | Chest and Secondary Incision Site | | | Chest Incision Only | | | | | | Overall | Initial | | | Revision | | | Overall | Initial | | | Revision | | | Overall | | | |
| | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | | | | |
| Saint Anthony Hospital | 0.00 | 0.00 | 29 | 5.17 | 6.90 | 58 | | | | | | | | | | | 0.47 | 0.94 | 212 | 0.00 | 0.00 | 4 | 0.46 | 0.93 | 0.38 | 0.38 | 263 | 10.00 | 10.00 | 10 | 0.73 | 0.73 | |
| Saint Clare Hospital | 0.00 | 0.00 | 10 | 2.86 | 11.43 | 35 | | | | | | | | | | | 0.53 | 1.05 | 190 | 0.00 | 0.00 | 8 | 0.51 | 1.01 | 0.45 | 0.45 | 224 | 0.00 | 0.00 | 8 | 0.43 | 0.43 | |
| Saint Elizabeth Hospital | 0.00 | 0.00 | 3 | 0.00 | 0.00 | 18 | | | | | | | | | | | 0.00 | 0.00 | 30 | 0.00 | 0.00 | 2 | 0.00 | 0.00 | 0.00 | 0.00 | 60 | 0.00 | 0.00 | 1 | 0.00 | 0.00 | |
| Saint Francis Hospital ¹⁰ | 3.03 | 3.03 | 33 | 6.25 | 8.75 | 80 | | | | | | | | | | | 0.00 | 0.00 | 80 | 33.33 | 33.33 | 3 | 1.20 | 1.20 | 0.00 | 0.00 | 92 | 0.00 | 0.00 | 3 | 0.00 | 0.00 | |
| Saint Joseph Medical Center | 0.69 | 0.92 | 434 | 3.85 | 7.14 | 182 | 0.83 | 1.65 | 242 | 0.00 | 0.00 | 25 | 0.75 | 1.50 | 0.54 | 0.54 | 186 | 0.55 | 0.82 | 366 | 0.00 | 0.00 | 22 | 0.52 | 0.77 | 0.43 | 1.07 | 467 | 0.00 | 0.00 | 30 | 0.40 | 1.01 |
| Samaritan Healthcare ¹⁶ | 0.00 | 0.00 | 8 | 5.88 | 17.65 | 17 | | | | | | | | | | | 1.54 | 1.54 | 65 | 0.00 | 0.00 | 6 | 1.41 | 1.41 | 0.00 | 0.00 | 142 | 0.00 | 0.00 | 9 | 0.00 | 0.00 | |
| Seattle Children's Hospital ¹¹ | 20.00 | 40.00 | 5 | 0.00 | 4.88 | 41 | - | - | 0 | - | - | 0 | - | - | 0.39 | 0.78 | 255 | 0.00 | 0.00 | 7 | - | - | 0 | 0.00 | 0.00 | 0.00 | 0.00 | 4 | 0.00 | 0.00 | 4 | 0.00 | 0.00 |
| Shriners Hospital for Children | | | | | | | | | | | | | | | | | 0.00 | 0.00 | 3 | - | - | 0 | 0.00 | 0.00 | - | - | 0 | - | - | 0 | - | - | |
| Skagit Valley Hospital ^{12, 23} | 3.85 | 3.85 | 26 | 2.67 | 2.67 | 75 | | | | | | | | | | | 1.75 | 2.63 | 114 | 33.33 | 33.33 | 3 | 2.56 | 3.42 | 0.55 | 0.55 | 182 | 0.00 | 0.00 | 18 | 0.50 | 0.50 | |
| Skyline Hospital | - | | 0 | - | - | 0 | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Sunnyside Community Hospital ²⁸ | 0.00 | 0.00 | 6 | 0.00 | 0.00 | 6 | | | | | | | | | | | 0.00 | 0.00 | 10 | - | - | 0 | 0.00 | 0.00 | 0.00 | 7.14 | 14 | 0.00 | 0.00 | 2 | 0.00 | 6.25 | |
| Swedish Ballard | 0.00 | 0.00 | 6 | 0.00 | 0.00 | 1 | | | | | | | | | | | 0.00 | 0.50 | 400 | 0.00 | 0.00 | 21 | 0.00 | 0.48 | 0.79 | 0.79 | 127 | 0.00 | 0.00 | 8 | 0.74 | 0.74 | |
| Swedish Cherry Hill | - | | 0 | - | - | 0 | 0.33 | 2.66 | 301 | 2.38 | 3.57 | 84 | 0.78 | 2.86 | 0.00 | 0.00 | 432 | | | | | | | | | | | | | | | | |
| Swedish Edmonds | 1.52 | 1.52 | 66 | 0.71 | 2.13 | 141 | | | | | | | | | | | 1.45 | 2.17 | 138 | 0.00 | 0.00 | 7 | 1.38 | 2.07 | 0.00 | 0.00 | 190 | 0.00 | 0.00 | 8 | 0.00 | 0.00 | |
| Swedish First Hill | 0.73 | 1.58 | 825 | 4.38 | 7.73 | 388 | | | | | | | | | | | 0.29 | 0.39 | 1020 | 1.08 | 1.62 | 185 | 0.41 | 0.58 | 0.46 | 0.61 | 1316 | 1.02 | 2.04 | 98 | 0.50 | 0.71 | |

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2013^{1, 2, 3}
October 8, 2014 download of January 2013 – December 2013 NHSN data

| Hospitals | Type of Surgical Procedure ⁴ | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|--|---|---------------|----------------------|------------------|---------------|----------------------|--|---------------|----------------------|------------------------|---------------|----------------------|------------------|---------------|----------------------|------------------------------|---------------|----------------------|--------------|---------------|----------------------|-------------------------------|---------------|----------------------|--------------|---------------|----------------------|--------------|---------------|---------|----|------|------|
| | Abdominal Hysterectomy | | | Colon Surgery | | | Coronary Artery Bypass Graft ⁵ | | | | | | Heart Surgery | | | Hip Replacement ⁵ | | | | | | Knee Replacement ⁵ | | | | | | | | | | | |
| | | | | | | | Chest and Secondary Incision Site | | | Chest Incision Only | | | | | | Overall | Initial | | | Revision | | | Overall | Initial | | | Revision | | | Overall | | | |
| | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | | | | |
| Swedish Issaquah | 1.60 | 3.20 | 125 | 1.54 | 7.69 | 65 | | | | | | | | | | | | 0.79 | 0.79 | 127 | 0.00 | 0.00 | 5 | 0.76 | 0.76 | 0.47 | 0.47 | 213 | 0.00 | 0.00 | 8 | 0.45 | 0.45 |
| Tacoma General Hospital / Allenmore Hospital | 0.00 | 0.00 | 387 | 3.57 | 4.46 | 224 | 0.42 | 0.42 | 237 | 0.00 | 0.00 | 6 | 0.41 | 0.41 | 0.93 | 0.93 | 108 | 1.69 | 1.69 | 236 | 0.00 | 0.00 | 12 | 1.61 | 1.61 | 0.00 | 0.33 | 307 | 7.14 | 7.14 | 28 | 0.60 | 0.90 |
| Three-Rivers Hospital | | | | 0.00 | 0.00 | 2 | | | | | | | | | | | | 0.00 | 0.00 | 8 | 0.00 | 0.00 | 1 | 0.00 | 0.00 | 0.00 | 0.00 | 9 | 0.00 | 0.00 | 5 | 0.00 | 0.00 |
| Toppenish Community Hospital | 0.00 | 0.00 | 7 | - | - | 0 | | | | | | | | | | | | 0.00 | 0.00 | 1 | - | - | 0 | 0.00 | 0.00 | 0.00 | 0.00 | 4 | - | - | 0 | 0.00 | 0.00 |
| Tri-State Memorial Hospital | 0.00 | 0.00 | 42 | 0.00 | 0.00 | 1 | | | | | | | | | | | | 0.00 | 0.00 | 89 | 0.00 | 0.00 | 10 | 0.00 | 0.00 | 0.00 | 0.58 | 172 | 0.00 | 0.00 | 18 | 0.00 | 0.53 |
| United General Hospital | | | | 0.00 | 0.00 | 10 | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| University of Washington Medical Center | 0.00 | 1.54 | 195 | 0.96 | 1.27 | 314 | 0.00 | 0.00 | 101 | 0.00 | 0.00 | 17 | 0.00 | 0.00 | 0.32 | 0.32 | 309 | 1.89 | 1.89 | 53 | 0.00 | 0.00 | 14 | 1.49 | 1.49 | 0.00 | 0.00 | 16 | 0.00 | 0.00 | 7 | 0.00 | 0.00 |
| Valley General Hospital | 0.00 | 0.00 | 1 | 0.00 | 0.00 | 36 | | | | | | | | | | | | 0.00 | 0.00 | 29 | 0.00 | 0.00 | 3 | 0.00 | 0.00 | 0.00 | 0.00 | 48 | 0.00 | 0.00 | 4 | 0.00 | 0.00 |
| Valley Hospital and Medical Center | 0.00 | 0.00 | 84 | 4.17 | 4.17 | 72 | | | | | | | | | | | | 1.07 | 1.60 | 187 | 0.00 | 0.00 | 19 | 0.97 | 1.46 | 0.00 | 0.00 | 306 | 0.00 | 0.00 | 2 | 0.00 | 0.00 |
| Valley Medical Center | 1.89 | 1.89 | 53 | 2.38 | 2.38 | 42 | | | | | | | | | | | | 1.66 | 1.90 | 422 | 1.96 | 1.96 | 51 | 1.69 | 1.90 | 0.98 | 0.98 | 613 | 0.00 | 0.00 | 47 | 0.91 | 0.91 |
| Virginia Mason Medical Center | 2.08 | 2.08 | 96 | 2.42 | 3.93 | 331 | 0.42 | 0.84 | 238 | 0.00 | 0.00 | 13 | 0.40 | 0.80 | 0.49 | 0.49 | 205 | 0.50 | 1.00 | 602 | 0.00 | 0.00 | 67 | 0.45 | 0.90 | 0.00 | 0.11 | 872 | 0.00 | 0.00 | 74 | 0.00 | 0.11 |
| Walla Walla General Hospital ²⁹ | 0.00 | 0.00 | 8 | 0.00 | 3.57 | 28 | | | | | | | | | | | | 0.00 | 0.00 | 25 | 0.00 | 0.00 | 2 | 0.00 | 0.00 | 2.56 | 2.56 | 39 | 0.00 | 0.00 | 2 | 2.44 | 2.44 |
| Wenatchee Valley Medical Center | 0.00 | 0.00 | 175 | - | - | 0 | | | | | | | | | | | | 0.00 | 0.00 | 44 | - | - | 0 | 0.00 | 0.00 | 0.00 | 0.00 | 69 | 0.00 | 0.00 | 1 | 0.00 | 0.00 |
| Whidbey General Hospital | 0.00 | 0.00 | 12 | 0.00 | 0.00 | 15 | | | | | | | | | | | | 0.00 | 2.04 | 49 | 0.00 | 0.00 | 2 | 0.00 | 1.96 | 0.00 | 1.47 | 68 | 0.00 | 0.00 | 3 | 0.00 | 1.41 |
| Whitman Hospital and Medical Center ^{17, 24} | - | - | 0 | 20.00 | 20.00 | 5 | | | | | | | | | | | | 0.00 | 6.67 | 15 | 0.00 | 0.00 | 2 | 0.00 | 5.88 | 0.00 | 0.00 | 33 | 0.00 | 0.00 | 1 | 0.00 | 0.00 |

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2013^{1, 2, 3}

October 8, 2014 download of January 2013 – December 2013 NHSN data

| Hospitals | Type of Surgical Procedure ⁴ | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|---|---|---------------|----------------------|------------------|---------------|----------------------|--|---------------|----------------------|------------------------|---------------|----------------------|------------------|---------------|----------------------|------------------------------|---------------|----------------------|--------------|---------------|----------------------|-------------------------------|---------------|----------------------|--------------|---------------|----------------------|--------------|---------------|------|----|------|------|
| | Abdominal Hysterectomy | | | Colon Surgery | | | Coronary Artery Bypass Graft ⁵ | | | | | | Heart Surgery | | | Hip Replacement ⁵ | | | | | | Knee Replacement ⁵ | | | | | | | | | | | |
| | | | | | | | Chest and Secondary Incision Site | | | Chest Incision Only | | Overall | | | | Initial | | Revision | | Overall | | Initial | | Revision | | Overall | | | | | | | |
| | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | No. of Procedures | Deep Rate | Total Rate | | | | |
| Willapa Harbor Hospital | | | | 0.00 | 0.00 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Yakima Regional Medical and Cardiac Center ²¹ | 0.00 | 0.00 | 23 | 3.03 | 4.55 | 66 | 0.00 | 0.67 | 150 | - | - | 0 | 0.00 | 0.67 | 1.49 | 1.49 | 67 | 0.00 | 0.00 | 45 | 11.11 | 11.11 | 9 | 1.85 | 1.85 | 0.00 | 0.00 | 123 | 0.00 | 0.00 | 15 | 0.00 | 0.00 |
| Yakima Valley Memorial Hospital ¹³ | 4.00 | 4.00 | 25 | 2.90 | 5.80 | 69 | | | | | | | | | | | | 0.46 | 0.93 | 216 | 0.00 | 0.00 | 9 | 0.44 | 0.89 | 0.57 | 0.85 | 353 | 0.00 | 6.67 | 15 | 0.54 | 1.09 |

Notes:

- All rates are a percent calculated as the number of surgical site infections (SSI) among all similar surgical procedures during the year (infections per 100 procedures). Red colored rate cells have statistically significant higher surgical infection rates than other hospitals, or (for those with an asterisk “*”) indicate a statistically high standardized infection ratio (SIR). See SIR description below. White rate cells have infection rates similar to other hospitals. Gold/yellow cells indicate a statistically low SIR. Light grey cells indicate hospitals do not report performing that type of surgical procedure. Visit our website for more information about our [statistical methods](#).

The annual number of procedures performed by a hospital is also provided as context to explain infection rates, and inform patients and providers in planning for elective surgery. In general, surgical teams which perform more of the same procedures have better outcomes.

- Depth of infection classification:

- The deep infection rate includes surgical wound infections occurring well below the skin and subcutaneous tissue, affecting muscles, organs, or bones (“deep incisional” and “organ/space” SSI per NHSN). Deep infections may be severe and can be difficult to treat often requiring antibiotics and a return visit to a hospital.
- The total infection rate includes deep infections as well as shallow surgical wound infections occurring within the skin and subcutaneous tissue (“superficial incisional” SSI per NHSN). Shallow infections can be more easily treated, often without hospitalization and may not require antibiotics. More information about the depth of [surgical site infections](#).
- Note – Rates for deep and total infections from the same procedure type use the same denominator (number of procedures performed) during the reporting period. See [box plots](#) for a graphical representation of the distribution of infection rates.

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2013^{1, 2, 3}

October 8, 2014 download of January 2013 – December 2013 NHSN data

3. The standardized infection ratio (SIR) is a summary statistic used by the CDC/NHSN to track healthcare associated infection prevention progress over time. The SIR compares reported HAI cases in a facility to “predicted” cases from historic baseline rates from the general U.S. population (national baseline for SSI is January 2006 – December 2008). The SIR for a facility is adjusted to account for factors that might cause infection rates to be higher or lower, such as hospital, surgery and patient characteristics. Washington State hospitals with surgical site infection SIRs significantly different than the national baseline are identified as either statistically high (red), or low (gold/yellow).
4. Visit our website for descriptions of these [types of surgical procedure categories](#).
5. Details of certain surgical procedure types are included in this table (coronary artery bypass graft procedures and joint replacement procedures). Hospitals are compared by their “overall” infection rates in these categories to compare hospitals more accurately. However, the more specific infection rate and procedure volume information in this table may be useful for patients and providers in planning for elective surgery.

Abdominal Hysterectomies

6. Grays Harbor Community Hospital – Infection rate is due to one deep surgical wound infection with a low number of procedures performed. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed. Additions were made to post-operative monitoring and patient follow-up after discharge.
7. Harborview Medical Center – Infection rate is due to two shallow surgical wound infections, with no indication of cross infection. The hospital reviewed both cases and reports that surgical care and infection prevention measures were followed. The hospital also reports enhanced infection surveillance due to a large outpatient clinic network.
8. Kennewick General Hospital – Infection rate is due to four surgical wound infections, two shallow and two deep, with no indication of cross-infection. The hospital’s standardized infection ratio (SIR), as calculated by CDC/NHSN, is also significantly higher than the national baseline. The hospital reviewed each case with their surgery committee and reports changing operating room work-flow and performed surgeon reeducation.
9. Providence Holy Family Hospital – Infection rate is due to five surgical wound infections, one shallow and four deep, with no indication of cross-infection. The hospital’s standardized infection ratio (SIR), as calculated by CDC/NHSN, is also significantly higher than the national baseline. The hospital investigated each case and has formed a multidisciplinary review board for infections, with administrative support. Surgical care processes have also been revised and operating room staff reeducated on infection prevention measures.
10. Saint Francis Hospital - Infection rate is due to only one deep surgical wound infection. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed.
11. Seattle Children’s Hospital – Infection rate is due to two surgical wound infections, one shallow and one deep, with no indication of cross infection and only five hysterectomy procedures performed during the year. The hospital operates on children with complex anatomical anomalies undergoing reconstructive surgeries. The hospital reviewed both cases and reports that surgical care and infection prevention measures were followed.
12. Skagit Valley Hospital – Infection rate is due to only one deep surgical wound infection. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed.
13. Yakima Valley Memorial Hospital – Infection rate is due to only one deep surgical wound infection. The case was reviewed and improvements were made to surgical care processes and infection prevention measures reinforced.

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2013^{1, 2, 3}

October 8, 2014 download of January 2013 – December 2013 NHSN data

Colon Surgeries

14. Central Washington Hospital – The hospital's standardized infection ratio (SIR), as calculated by CDC/NHSN, is significantly higher than the national baseline, with two times the predicted number of infections occurring throughout the year. The hospital reviewed each case and has made improvements to their operative infection prevention practices. The hospital also reports enhanced infection surveillance from access to an outpatient clinic network.
15. Pullman Regional Hospital – Infection rate is due to two deep surgical wound infections, with no indication of cross-infection and a low number of colon procedures performed during the year. The hospital investigated both cases and identified a potential issue with a surgical closure device in both cases and has since discontinued the use of that device brand. Surgical site prevention practices have been reinforced and staff reeducated.
16. Samaritan Healthcare – Infection rate is due to three surgical wound infections, two shallow and one deep, with no indication of cross-infection and a low number of colon procedures performed during the year. The hospital reviewed each case and reports that surgical care and infection prevention measures were followed.
17. Whitman Hospital and Medical Center – Infection rate is due to one deep surgical wound infection out of only five colon procedures performed during the year. The hospital reports all appropriate infection prevention measures have been in place and continue to be enforced.

Coronary Artery Bypass Graft Surgeries

For coronary artery bypass graft surgeries, hospitals are compared by their “overall” infection rates, to compare hospitals more accurately. Details of coronary artery bypass graft surgeries with secondary incision sites (CBGB) or chest incision only (CBGC) may be useful for patients and providers in planning for elective surgery and for comparison to other reports using these NHSN categories.

18. Deaconess Medical Center – Infection rate is due to three deep surgical wound infections with no indication of cross-infection. The hospital investigated each case and reports increasing surgical staff, with reeducation and reinforcing infection prevention measures. Changes were also made to artery donor site incision practices.
19. Northwest Hospital – Infection rate is due to only one deep surgical wound infection with a low number of coronary artery procedures performed during the year. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed. Recent updates in cardiac surgery infection prevention guidelines have also been added to their routine pre-operative measures.

Heart (Cardiac) Surgeries

20. PeaceHealth Saint Joseph Medical Center – Infection rate is due to two deep surgical wound infections with no indication of cross-infection. The hospital reviewed the cases with a multi-disciplinary team and reports that surgical care and infection prevention measures were followed.
21. Yakima Regional Medical and Cardiac Center – Infection rate is due to only one deep surgical wound infection with a low number of heart surgeries performed during the year. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed.

Table of Surgical Site Infection Rates, by Type of Procedure and Depth of Infection, 2013^{1, 2, 3}

October 8, 2014 download of January 2013 – December 2013 NHSN data

Hip Replacement Surgeries

For hip replacement surgeries, hospitals are compared by their “overall” infection rates, to compare hospitals more accurately. Details of initial (“primary” per NHSN) and repeat or revision hip prosthesis surgeries may be useful for patients and providers in planning for elective surgery.

22. Jefferson General Hospital – Infection rate is due to four surgical wound infections, two shallow and two deep, with no indication of cross-infection and a low number of hip replacement surgeries during the year. The hospital reviewed each case and has reinforced surgical care measures, made improvements to their operative infection prevention practices and has formed a multidisciplinary team to review infections.
23. Skagit Valley Hospital – Infection rate is due to three deep surgical wound infections throughout the year. The hospital reports that surgical care measures were followed. The hospital is working to improve processes and communication between the surgeons, surgical teams, and patients to prevent and investigate infections.
24. Whitman Hospital and Medical Center – Infection rate is due to one shallow surgical wound infection with a low number of hip procedures performed during the year. The hospital reports all appropriate infection prevention measures have been in place and continue to be enforced.


Knee Replacement Surgeries

For knee replacement surgeries, hospitals are compared by their “overall” infection rates, to compare hospitals more accurately. Details of initial (“primary” per NHSN) and repeat or revision knee prosthesis surgeries may be useful for patients and providers in planning for elective surgery.

25. Harborview Medical Center – Infection rate is due to only one deep surgical wound infection out of only five knee replacement procedures performed during the year. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed. The hospital does not perform elective joint replacement surgery. The patients that receive these procedures are admitted for traumatic injuries.
26. Jefferson General Hospital – Infection rate is due to only one shallow surgical wound infection with a low number of knee replacement surgeries during the year. The hospital reviewed each case and has reinforced surgical care measures, made improvements to their operative infection prevention practices and has formed a multidisciplinary team to review infections.
27. Kennewick General Hospital – Infection rate is due to two deep surgical wound infections with no indication of cross-infection. The hospital reviewed each case with their surgery committee and reports changing operating room work-flow and that surgical care and infection prevention measures were followed.
28. Sunnyside Community Hospital – Infection rate is due to only one shallow surgical wound infection and a very low number of knee replacement surgeries during the year. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed.
29. Walla Walla General Hospital – Infection rate is due to only one deep surgical wound infection. The hospital reviewed the cases and reports that surgical care and infection prevention measures were followed. The hospital also reports changes to preoperative procedures including pre-surgery patient preparation and education.

Box Plot of Surgical Site Infection Rates, by Procedure Type and Infection Depth, 2013¹

October 8, 2014 download of January 2013 – December 2013 NHSN data



Notes:

1. All rates are a percent calculated as the number of surgical site infections (SSI) among all similar surgical procedures (infections per 100 procedures). Hospitals identified in the plots below, have statistically significantly higher surgical infection rates than other hospitals. Visit our website for more information about our [statistical methods](#).

The annual number of procedures performed by a hospital (in the table and map format of the [annual SSI reports](#)) provides more context to explain certain infection rates and inform patients and providers in planning for elective surgery. In general, surgical teams which perform more of the same procedures have better outcomes.

2. The distribution of hospital infection rates for all six Washington State reportable surgery types is included in this box plot. Below are more detailed boxplots grouping these procedures into three broad categories ([abdominal surgeries](#), [heart surgeries](#), and [joint replacements](#)). Visit our website for descriptions of these [types of surgeries](#).
3. Depth of [surgical site infection](#) classification:
 - a. Deep infection rates include surgical wound infections occurring well below the skin and subcutaneous tissue, affecting muscles, organs, or bones (“deep incisional” and “organ/space” SSI per NHSN). Deep infections may be severe and difficult to treat, often requiring antibiotics and additional hospital care.
 - b. Total infection rates include deep infections as well as shallow surgical wound infections occurring within the skin and subcutaneous tissue (“superficial incisional” SSI per NHSN). Shallow infections can be more easily treated without hospitalization and may not require antibiotics.
 - c. Rates for deep and total infections from the same procedure type use the same denominator (number of procedures performed) during the reporting period.

Box Plot of Surgical Site Infection Rates, by Procedure Type and Infection Depth, 2013 ¹


October 8, 2014 download of January 2013 – December 2013 NHSN data

4. Rates shown here are for the surgical procedure overall. Details of certain surgical procedure types are included in the [SSI table](#) (coronary artery bypass graft procedures and joint replacement procedures). Hospitals are compared by their “overall” infection rates in these categories to compare hospitals more accurately. However, the more specific infection rate and procedure volume information in the table may be useful for patients and providers in planning for elective surgery.

Box Plot of Surgical Site Infection Rates, by Procedure Type and Infection Depth, 2013 ¹

October 8, 2014 download of January 2013 – December 2013 NHSN data

Abdominal Surgeries



Abdominal Hysterectomies

5. Grays Harbor Community Hospital – Infection rate is due to one deep surgical wound infection with a low number of procedures performed. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed. Additions were made to post-operative monitoring and patient follow-up after discharge.
6. Harborview Medical Center – Infection rate is due to two shallow surgical wound infections, with no indication of cross infection. The hospital reviewed both cases and reports that surgical care and infection prevention measures were followed. The hospital also reports enhanced infection surveillance due to a large outpatient clinic network.
7. Kennewick General Hospital – Infection rate is due to four surgical wound infections, two shallow and two deep, with no indication of cross-infection. The hospital's standardized infection ratio (SIR), as calculated by CDC/NHSN, is also significantly higher than the national baseline. The hospital reviewed each case with their surgery committee and reports changing operating room work-flow and performed surgeon reeducation.
8. Providence Holy Family Hospital – Infection rate is due to five surgical wound infections, one shallow and four deep, with no indication of cross-infection. The hospital's standardized infection ratio (SIR), as calculated by CDC/NHSN, is also significantly higher than the national baseline. The hospital investigated each case and has formed a multidisciplinary review board for infections, with administrative support. Surgical care processes have also been revised and operating room staff reeducated on infection prevention measures.
9. Saint Francis Hospital - Infection rate is due to only one deep surgical wound infection. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed.
10. Seattle Children's Hospital – Infection rate is due to two surgical wound infections, one shallow and one deep, with no indication of cross infection and only five hysterectomy procedures performed during the year. The hospital operates on children with complex anatomical anomalies undergoing reconstructive surgeries. The hospital reviewed both cases and reports that surgical care and infection prevention measures were followed.
11. Skagit Valley Hospital – Infection rate is due to only one deep surgical wound infection. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed.

Box Plot of Surgical Site Infection Rates, by Procedure Type and Infection Depth, 2013 ¹

October 8, 2014 download of January 2013 – December 2013 NHSN data

12. Yakima Valley Memorial Hospital – Infection rate is due to only one deep surgical wound infection. The case was reviewed and improvements were made to surgical care processes and infection prevention measures reinforced.


Colon Surgeries

13. Central Washington Hospital (*not pictured*) – The hospital's standardized infection ratio (SIR), as calculated by CDC/NHSN, is significantly higher than the national baseline, with two times the predicted number of infections occurring throughout the year. The hospital reviewed each case and has made improvements to their operative infection prevention practices. The hospital also reports enhanced infection surveillance from access to an outpatient clinic network.
14. Pullman Regional Hospital – Infection rate is due to two deep surgical wound infections, with no indication of cross-infection and a low number of colon procedures performed during the year. The hospital investigated both cases and identified a potential issue with a surgical closure device in both cases and has since discontinued the use of that device brand. Surgical site prevention practices have been reinforced and staff reeducated.
15. Samaritan Healthcare – Infection rate is due to three surgical wound infections, two shallow and one deep, with no indication of cross-infection and a low number of colon procedures performed during the year. The hospital reviewed each case and reports that surgical care and infection prevention measures were followed.
16. Whitman Hospital and Medical Center – Infection rate is due to one deep surgical wound infection out of only five colon procedures performed during the year. The hospital reports all appropriate infection prevention measures have been in place and continue to be enforced.

Box Plot of Surgical Site Infection Rates, by Procedure Type and Infection Depth, 2013 ¹

October 8, 2014 download of January 2013 – December 2013 NHSN data

Heart Surgeries



Coronary Artery Bypass Graft Surgeries

Rates shown here are for coronary artery bypass graft surgeries overall. Details of coronary artery bypass graft surgeries with secondary incision sites (CBGB) or chest incision only (CBGC) are included in the [2013 SSI table](#). Hospitals are compared by their “overall” infection rates in these categories to compare hospitals more accurately.

17. Deaconess Medical Center – Infection rate is due to three deep surgical wound infections with no indication of cross-infection. The hospital investigated each case and reports increasing surgical staff, with reeducation and reinforcing infection prevention measures. Changes were also made to artery donor site incision practices.
18. Northwest Hospital – Infection rate is due to only one deep surgical wound infection with a low number of coronary artery procedures performed during the year. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed. Recent updates in cardiac surgery infection prevention guidelines have also been added to their routine pre-operative measures.


Heart (Cardiac) Surgery

19. PeaceHealth Saint Joseph Medical Center – Infection rate is due to two deep surgical wound infections with no indication of cross-infection. The hospital reviewed the cases with a multi-disciplinary team and reports that surgical care and infection prevention measures were followed.
20. Yakima Regional Medical and Cardiac Center – Infection rate is due to only one deep surgical wound infection with a low number of heart surgeries performed during the year. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed.

Box Plot of Surgical Site Infection Rates, by Procedure Type and Infection Depth, 2013¹

October 8, 2014 download of January 2013 – December 2013 NHSN data

Joint Replacements



Hip Replacement

Rates shown here are for all hip replacement surgeries overall. Details of initial (primary) and repeat or revision procedures are included in the [2013 SSI table](#). Hospitals are compared by their “overall” infection rates in these categories to compare hospitals more accurately.

21. Jefferson General Hospital – Infection rate is due to four surgical wound infections, two shallow and two deep, with no indication of cross-infection and a low number of hip replacement surgeries during the year. The hospital reviewed each case and has reinforced surgical care measures, made improvements to their operative infection prevention practices and has formed a multidisciplinary team to review infections.
22. Skagit Valley Hospital – Infection rate is due to three deep surgical wound infections throughout the year. The hospital reports that surgical care measures were followed. The hospital is working to improve processes and communication between the surgeons, surgical teams, and patients to prevent and investigate infections.
23. Whitman Hospital and Medical Center – Infection rate is due to one shallow surgical wound infection with a low number of hip procedures performed during the year. The hospital reports all appropriate infection prevention measures have been in place and continue to be enforced.

Knee Replacement

Rates shown here are for all knee replacement surgeries overall. Details of initial (primary) and repeat or revision procedures are included in the [2013 SSI table](#). Hospitals are compared by their “overall” infection rates in these categories to compare hospitals more accurately.

24. Harborview Medical Center – Infection rate is due to only one deep surgical wound infection out of only five knee replacement procedures performed during the year. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed. The hospital does not perform elective joint replacement surgery. The patients that receive these procedures are admitted for traumatic injuries.

Box Plot of Surgical Site Infection Rates, by Procedure Type and Infection Depth, 2013 ¹

October 8, 2014 download of January 2013 – December 2013 NHSN data

25. Jefferson General Hospital – Infection rate is due to only one shallow surgical wound infection with a low number of knee replacement surgeries during the year. The hospital reviewed each case and has reinforced surgical care measures, made improvements to their operative infection prevention practices and has formed a multidisciplinary team to review infections.
26. Kennewick General Hospital – Infection rate is due to two deep surgical wound infections with no indication of cross-infection. The hospital reviewed each case with their surgery committee and reports changing operating room workflow and that surgical care and infection prevention measures were followed.
27. Sunnyside Community Hospital – Infection rate is due to only one shallow surgical wound infection and a very low number of knee replacement surgeries during the year. The hospital reviewed the case and reports that surgical care and infection prevention measures were followed.
28. Walla Walla General Hospital – Infection rate is due to only one deep surgical wound infection. The hospital reviewed the cases and reports that surgical care and infection prevention measures were followed. The hospital also reports changes to preoperative procedures including pre-surgery patient preparation and education.