

VACCINE	ELIGIBILITY
DTaP / DT	Children from 6 weeks of age up to the 7th birthday
DTaP/IPV/Hep B Combination (Pediatrix [®])	<p>Children from 2 months of age up to the 7th birthday:</p> <ul style="list-style-type: none"> Indicated for the primary doses of DTaP, IPV, and Hep B series at 2, 4 and 6 months of age. Pediatrix may be used for catch up vaccination for children up to 7 years of age who have not completed the primary series. <p>Considerations: If Pediatrix[®] is not available, single injection vaccines may replace any dose in the vaccination series. Pediatrix[®] should not be given to infants less than 6 weeks of age or any child 7 years of age or older. Pediatrix[®] does not replace the hepatitis B birth dose. ACIP indicates that it is acceptable for children to receive 4 doses of hepatitis B vaccine.</p>
DTaP-IPV/Hib Combination (Pentacel [®])	<p>Children from 6 weeks of age up to the 5th birthday:</p> <ul style="list-style-type: none"> Indicated for the 4 dose DTaP, IPV, and Hib series at 2, 4, 6 and 15 – 18 months of age. <p>Considerations: If Pentacel[®] is not available, single injection vaccines may replace any dose in the vaccination series. DTaP-IPV/Hib is not licensed for children aged ≥ 5 years or for the 5th doses of the DTaP series. DTaP-IPV/Hib inadvertently administered to children aged ≥ 5 years should be counted as a valid dose.</p>
DTaP-IPV (KINRIX [®])	<p>Children from 4 years through 6 years of age:</p> <ul style="list-style-type: none"> Indicated for the booster dose (5th dose in the DTaP series and the 4th dose in the IPV series)
Hepatitis A	<p>Children from 12 months of age up to the 19th birthday:</p> <ul style="list-style-type: none"> All children should receive hepatitis A vaccine at 1 year of age (i.e., 12-23 months). 2 doses of hepatitis A vaccine are required. The minimum interval between the first and second dose is 6 months.
Hepatitis B	<p>Children from birth up to the 19th birthday:</p> <p>Targeted Groups:</p> <ul style="list-style-type: none"> Children born on or after 11/22/91 Adolescents 11 & 12 years of age <p>Targeted High Risk Groups:</p> <ul style="list-style-type: none"> Children from birth up to the 20th birthday who meet the high risk criteria: <ol style="list-style-type: none"> Children born after 10/1/87 to 1st generation immigrant women from countries of high or intermediate hepatitis B virus endemicity Persons with occupational risk Clients in institutions for the developmentally disabled Hemodialysis patients Recipients of certain blood products Household contacts/sexual partners of HBV carriers Adoptees from countries where HBV is endemic International travelers Injecting drug users Sexually active homosexual and bisexual men Sexually active heterosexual men and women Inmates of long-term correctional facilities

VACCINE	ELIGIBILITY
Hep B – Hib (COMVAX®)	<p>Children from 6 weeks of age up to 15 months of age:</p> <ul style="list-style-type: none"> • Do not administer to infants less than 6 weeks of age. • Must not be used for doses at birth or 1 month of age for a child on a 0, 1, 6 month hepatitis B vaccine schedule. • Licensed for use at 2, 4, and 15-18 months of age.
Hib	<p>Children from 6 weeks of age up to the 5th birthday</p>
HPV -- Gardasil (quadravalent – types 6, 11, 16 and 18)	<p>HPV vaccine is routinely recommended:</p> <ul style="list-style-type: none"> • For adolescent females, 11 through 12 years of age (3 dose series). <p>Catch-up vaccination:</p> <ul style="list-style-type: none"> • Vaccination is recommended for females 13-18 years of age who have not been previously vaccinated or who have not completed the full series. <p>Other permitted vaccination:</p> <ul style="list-style-type: none"> • Adolescent females from 9 years of age up to the 19th birthday may be vaccinated. • Adolescent males from 9 years of age up to the 19th birthday may be vaccinated.
HPV -- Cervarix (bivalent – types 16 and 18)	<p>HPV vaccine is routinely recommended:</p> <ul style="list-style-type: none"> • For adolescent females, 11 through 12 years of age (3 dose series). <p>Catch-up vaccination:</p> <ul style="list-style-type: none"> • Vaccination is recommended for females 13-18 years of age who have not been previously vaccinated or who have not completed the full series. <p>Considerations:</p> <ul style="list-style-type: none"> • Adolescent females from 10 years of age up to the 19th birthday may be vaccinated. • Not approved for use in males.
IPV (Polio)	<p>Children from 6 weeks of age up to the 19th birthday</p>

VACCINE	ELIGIBILITY
Meningococcal (MCV-4)	<p>Children 11 up to the 19th birthday and children 2 – up to the 19th birthday at high risk:*</p> <p>Adolescents age 11 up to the 19th birthday:</p> <p>A single dose of meningococcal vaccine is recommended at</p> <ul style="list-style-type: none"> • 11 through 12 years of age for all adolescents. <p>A second dose (booster) of meningococcal vaccine is recommended at:</p> <ul style="list-style-type: none"> • Age 16 if the first dose was administered at 11-12 years of age. • Age 16 – 18 if the first dose was administered at 13 – 15 years of age. • <i>If</i> the first dose is administered at or after age 16, no booster dose is required. <p>*Children age 2 up to the 19th birthday meeting the following high risk criteria:</p> <p>A one dose primary series is recommended for:</p> <ul style="list-style-type: none"> • Children who travel to or reside in countries in which N. meningitidis is hyperendemic or epidemic, particularly if contact with the local population will be prolonged <p>A two dose primary series is recommended as follows:</p> <ul style="list-style-type: none"> • Children who have terminal complement component deficiencies • Children who have anatomic or functional asplenia • Children previously vaccinated with MPSV4 who remain at increased risk for meningococcal disease • Adolescents 11 years up to the 19th birthday infected with human immunodeficiency virus (HIV) <p>In addition to the primary series, a booster is recommended for children at high risk. The number and spacing of these boosters varies. Please see the ACIP recommendation or ACIP Recommended Schedule.</p> <p>http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6003a3.htm?s_cid=mm6003a3_e%0d%0a</p>
MMR	<p>Children from 12 months of age up to the 19th birthday:</p> <p><u>First dose:</u>*</p> <ul style="list-style-type: none"> • All persons 12 months of age up to the 19th birthday • Students entering college who were born in or after 1957 <p><u>Second dose:</u>*</p> <ul style="list-style-type: none"> • All children at 4 to 6 years of age • Any child less than 19 years of age who has not previously received a second dose • Students entering college who were born in or after 1957 who have not yet received a second dose <p>*Please consult with the state immunization program for appropriate use of MMR vaccine during measles outbreaks.</p>

VACCINE	ELIGIBILITY
MMRV (mumps, measles, rubella, varicella combination vaccine) (ProQuad®)	<p>Children from 12 months of age up to 12 years: Administer MMRV when MMR and varicella vaccine is indicated for the first or second dose as follows:</p> <ul style="list-style-type: none"> • Ages 12 – 47 months -- first dose of measles, mumps, rubella, and varicella vaccines: either measles, mumps, and rubella (MMR) vaccine and varicella vaccine or MMRV vaccine may be used. CDC recommends that MMR vaccine and varicella vaccine should be administered for the first dose in this age group. • Second dose of measles, mumps, rubella, and varicella vaccines at any age (15 months--12 years) or the first dose at ≥48 months of age, use of MMRV vaccine generally is preferred over separate injections of its equivalent component vaccines (i.e., MMR vaccine and varicella vaccine). <p>Considerations: MMRV (ProQuad™) must be used within 30 minutes of reconstitution. MMRV (ProQuad™) must be stored frozen at or below -15° C. (5° F.). Providers must be certified for MMRV. Contact your local health department regarding certification requirements. Providers currently certified for varicella will be considered certified for MMRV.</p>
Pneumococcal Conjugate (PCV)	<p>Children 2 months of age up to the 5th birthday:</p> <ul style="list-style-type: none"> • All children 2 months of age up to the 5th birthday at 2, 4, 6 and 12-15 months of age (although the vaccine can be given as early as 6 weeks of age) <p>Children aged 60-71 months of age with underlying medical conditions that increase their risk for pneumococcal disease or complications may receive a booster dose of this vaccine.</p> <p>See MMWR for ACIP Recommendations for other permitted vaccination: http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5909a2.htm?s_cid=mm5909a2_e</p>
Pneumococcal Polysaccharide	<p>Children from 2 years of age up to the 19th birthday who meet the following <u>high-risk</u> criteria:</p> <ul style="list-style-type: none"> • Asplenia • Sickle cell disease • Nephrotic syndrome • Cerebrospinal fluid leaks • Immunosuppression • Living in environments or social settings with an identified increased risk of pneumococcal disease or its complications
Rotavirus	<p>Infants aged 6 weeks through 8 months:</p> <ul style="list-style-type: none"> ▪ Rotateq® (RV5) – 3 dose series (2, 4 and 6 months of age) ▪ Rotarix® (RV1) – 2 dose series (2 and 4 months of age) ▪ 1st dose of rotavirus vaccine 6 -14 weeks 6 days (maximum age is 14 weeks 6 days) ▪ Minimum interval between doses: 4 weeks. <p>Considerations: Vaccination should not be initiated for infants age 15 weeks or older; minimum interval 4 weeks; all doses administered by 8 calendar months of age; use same product if possible – if any dose in the series is Rotateq® or unknown, use 3 dose series.</p>
Td	<p>Children from 7 years of age up to the 19th birthday for whom Tdap is contraindicated or unavailable</p>

VACCINE	ELIGIBILITY
Tdap	<p>Children from 11 years of age up to the 19th birthday: Tdap vaccine is routinely recommended:</p> <ul style="list-style-type: none"> • For adolescents at 11 through 12 years of age (a single dose). • Tdap may be used regardless of the interval since the last tetanus or diphtheria-toxoid containing vaccine. <p>Administration of Tdap for adolescents in special circumstances:</p> <ul style="list-style-type: none"> • Tdap is indicated but not available: Td can be administered if the last DTP/DTaP/DT/Td vaccine was equal to or greater than 10 years earlier. • Pertussis outbreaks and other setting with increased risk from pertussis: Routine Tdap vaccination recommendations for adolescents should be used. • Tetanus Prophylaxis in Wound Management: A single dose of Tdap instead of Td if they have not previously received Tdap. • No History of DTP/DTaP/Td/Tdap Vaccination: A single Tdap dose, followed by a dose of Td ≥4 weeks after the Tdap dose and a second dose of Td ≥6 months after the Td dose. Tdap may substitute for any one of the 3 Td doses in the series. • Pregnancy: If otherwise indicated, consider a single dose of Tdap for adolescents immediately after delivery. <p>Catch-up vaccination using state-supplied Tdap vaccine is allowed for children 7 through 18 years of age who have:</p> <ul style="list-style-type: none"> • Received tetanus and diphtheria containing vaccines (DT or Td) instead of DTP/DTaP for some or all doses of the childhood series. • Received fewer than 5 doses of DTP/DTaP or 4 doses if the fourth dose was administered at age 4 years or older. • Never been vaccinated against tetanus, diphtheria, or pertussis (no doses of pediatric DTP/DTaP/DT or Td). <p>The preferred schedule is a single Tdap dose, followed by a dose of Td four weeks after the first dose and a second dose of Td 6-12 months later. If not administered as the first dose, Tdap can be substituted for any of the other Td doses in the series.</p> <p>Regarding the off label use of Tdap: Tdap vaccines are not licensed for children less than 10 years of age. Use of Tdap vaccine for children less than 10 years of age (Boostrix) or less than 11 years of age (Adacel) is considered off-label use. In the event a provider makes a clinical decision to use Tdap vaccine "off-label," to vaccinate a 7 - 9 year old, state-supplied vaccine may be used. The liability for off-label use of any vaccine (or other medication) is the responsibility of the provider exercising his or her clinical judgment in making the decision to do so.</p>

VACCINE	ELIGIBILITY
Varicella (Chickenpox)	<p>Children at least 12 months of age up to the 19th birthday who do not have evidence of varicella immunity.</p> <p><u>First dose:</u></p> <ul style="list-style-type: none"> Children 12-15 months of age. Children less than 19 years of age who do not have evidence of varicella immunity, and have not received the first dose <p><u>Second dose:</u></p> <ul style="list-style-type: none"> Children 4-6 years of age Children less than 19 years of age who do not have evidence of varicella immunity, and have not received two doses of varicella vaccine. <p>Considerations:</p> <ul style="list-style-type: none"> Providers must be certified for MMRV. Contact your local health department regarding certification requirements.

VACCINE	ELIGIBILITY										
Seasonal Influenza (2011-2012)	<p>All children aged 6 months up to the 19th birthday:</p> <table border="1"> <thead> <tr> <th>Flu Vaccine</th> <th>Age Indications for Use of State-Supplied Vaccine</th> </tr> </thead> <tbody> <tr> <td>Sanofi Pasteur Fluzone[®] Preservative free, 0.25 mL Single Dose Syringe</td> <td>Children aged six months up to the third birthday.</td> </tr> <tr> <td>Novartis Fluvirin[®] Preservative free, 0.5 mL Single Dose Syringe</td> <td>Prioritize the use of this vaccine for pregnant adolescents less than age 19 years. If supplies are sufficient to vaccinate this group, this presentation may be used for children aged four years up to the nineteenth birthday who meet the criteria for the use of state-supplied vaccine.</td> </tr> <tr> <td>Sanofi Pasteur Fluzone[®] 5.0 mL Multi Dose Vial</td> <td>Children aged three years up to the nineteenth birthday.</td> </tr> <tr> <td>MedImmune FluMist[®] Preservative free , Single Dose Nasal Spray</td> <td>Children aged two years up to the nineteenth birthday.</td> </tr> </tbody> </table> <p>*Children less than 19 years of age are considered high risk if they meet the following criteria:</p> <ul style="list-style-type: none"> Chronic illness, i.e. chronic pulmonary or cardiovascular conditions, metabolic diseases, renal dysfunction, hemoglobinopathies. Conditions that compromise respiratory function or the handling of respiratory secretions or can increase the risk of aspiration. Children receiving chronic aspirin therapy. Children receiving immunosuppression therapy. 	Flu Vaccine	Age Indications for Use of State-Supplied Vaccine	Sanofi Pasteur Fluzone [®] Preservative free, 0.25 mL Single Dose Syringe	Children aged six months up to the third birthday.	Novartis Fluvirin [®] Preservative free, 0.5 mL Single Dose Syringe	Prioritize the use of this vaccine for pregnant adolescents less than age 19 years. If supplies are sufficient to vaccinate this group, this presentation may be used for children aged four years up to the nineteenth birthday who meet the criteria for the use of state-supplied vaccine.	Sanofi Pasteur Fluzone [®] 5.0 mL Multi Dose Vial	Children aged three years up to the nineteenth birthday.	MedImmune FluMist [®] Preservative free , Single Dose Nasal Spray	Children aged two years up to the nineteenth birthday.
Flu Vaccine	Age Indications for Use of State-Supplied Vaccine										
Sanofi Pasteur Fluzone [®] Preservative free, 0.25 mL Single Dose Syringe	Children aged six months up to the third birthday.										
Novartis Fluvirin [®] Preservative free, 0.5 mL Single Dose Syringe	Prioritize the use of this vaccine for pregnant adolescents less than age 19 years. If supplies are sufficient to vaccinate this group, this presentation may be used for children aged four years up to the nineteenth birthday who meet the criteria for the use of state-supplied vaccine.										
Sanofi Pasteur Fluzone [®] 5.0 mL Multi Dose Vial	Children aged three years up to the nineteenth birthday.										
MedImmune FluMist [®] Preservative free , Single Dose Nasal Spray	Children aged two years up to the nineteenth birthday.										

For questions or comments, contact the Immunization and Child Profile Office at 360-236-3595.