

*Behavioral Risk  
Factor Surveillance System*

*State Prevalence Report  
of Risk Factors for  
WASHINGTON Statewide  
2001*



T A B L E O F C O N T E N T S

TABLE	PAGE
STATE PREVALENCE REPORT OF RISK FACTORS FOR WASHINGTON STATEWIDE	2001
HEALTH INSURANCE: HAVE NO HEALTH CARE COVERAGE, AGES 18-64 .....	3
HEALTH CARE ACCESS: NO SPECIFIC SOURCE OF ON-GOING HEATH CARE, AGES 18 AND OVER .....	4
COLORECTAL SCREENING: NO FOBT IN PAST 2 YRS, AGE 50 AND OVER .....	5
SIGMOIDOSCOPY: NEVER HAD A SIGMOIDOSCOPY/COLONSCOPY, AGE 50 AND OVER .....	6
DIABETES: EVER TOLD YOU HAVE DIABETES BY A DOCTOR .....	7
CHOLESTEROL AWARENESS: NO CHOLESTEROL CHECK IN PAST 5 YEARS, AGE 18 AND OVER .....	8
FLU IMMUNIZATION: NO FLU SHOT IN PAST 12 MONTHS, AGE 65 AND OVER .....	9
PNEUMONIA VACCINATION: DID NOT HAVE A PNEUMONIA VACCINATION, AGE 65 AND OVER .....	10
OBESITY: OBESITY DEFINED BY BODY MASS INDEX, AGE 18 AND OVER .....	11
LEISURE TIME PHYSICAL ACTIVITY: NO PHYSICAL ACTIVITY IN PAST THIRTY DAYS, AGE 18 AND OVER .....	12
TOBACCO USE: CURRENT SMOKER, AGE 18 AND OVER .....	13
ALCOHOL CONSUMPTION: BINGE DRINKING DURING THE PAST THIRTY DAYS, AGE 18 AND OVER .....	14

NOTE :

THIS SECTION OF THE TABULATIONS CONTAINS PREVALENCE ESTIMATES FOR HEALTH BEHAVIORS AND PREVENTIVE HEALTH MEASURES BY RACE AND SEX.

FOR THESE TABLES, THE RACE GROUP 'NON-HISPANIC WHITES' IS DEFINED AS ALL RESPONDENTS WHO INDICATED THEY WERE WHITE AND NOT OF HISPANIC ORIGIN. THE GROUP 'HISPANIC OR NON-WHITES' IS DEFINED AS ALL RACES EXCLUDING NON-HISPANIC WHITE RESPONDENTS.

2001 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM  
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : HEALTH INSURANCE: HAVE NO HEALTH CARE COVERAGE: AGES 18-64 (HP 2010 OBJ. 1.1)

	TOTAL *****	MALES *****	FEMALES *****	HISP./MULTIRACE NON-WHITE *****	NON-HISPANIC WHITE *****
SAMPLE SIZE	3454	1489	1965	517	2892
PREVALENCE	11.6%	12.2%	10.9%	14.8%	10.9%
STANDARD ERROR	0.7	1.0	0.9	1.9	0.7
95% C.I.	( 10.2, 12.9)	( 10.3, 14.1)	( 9.0, 12.7)	( 11.1, 18.5)	( 9.5, 12.3)

	HISP./MULTIRACE NON-WHITE MALES	NON-HISPANIC WHITE MALES	HISP./MULTIRACE NON-WHITE FEMALES	NON-HISPANIC WHITE FEMALES
SAMPLE SIZE	242	1228	275	1664
PREVALENCE	16.2%	11.4%	13.0%	10.4%
STANDARD ERROR	2.9	1.0	2.2	1.0
95% C.I.	( 10.6, 21.8)	( 9.4, 13.5)	( 8.8, 17.3)	( 8.3, 12.4)

	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
<b>MALES/FEMALES</b> *****						
TOTAL	1.3%	1.4	( 0.0, 4.0)	0.96	1.12	( 0.89, 1.41)
HISP./MULTIRACE NON-WH	3.2%	3.6	( 0.0,10.2)	0.88	1.24	( 0.77, 2.02)
NON-HISPANIC WHITES	1.0%	1.5	( 0.0, 3.9)	0.70	1.10	( 0.84, 1.43)
<b>HISP./MULTIRACE NON-WHITES/NON-HISPANIC WHITES</b> *****						
TOTAL	3.9%	2.0	( 0.0, 7.8)	1.94	1.36	( 1.00, 1.85)
MALES	4.8%	3.1	( 0.0,10.8)	1.57	1.42	( 0.92, 2.20)
FEMALES	2.6%	2.4	( 0.0, 7.4)	1.09	1.25	( 0.84, 1.88)

\*DIFFERENCE IS SIGNIFICANT AT ALPHA = .05  
\*\*1.0 IS NOT IN THE CONFIDENCE INTERVAL

2001 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM  
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : NO SPECIFIC SOURCE OF ON-GOING HEALTH CARE: AGE 18 & OVER (HP 2010 OBJ. 1.4c)

	TOTAL *****	MALES *****	FEMALES *****	HISP./MULTIRACE NON-WHITE *****	NON-HISPANIC WHITE *****
SAMPLE SIZE	4194	1751	2443	574	3568
PREVALENCE	20.6%	26.5%	14.9%	26.1%	19.6%
STANDARD ERROR	0.8	1.3	0.9	2.4	0.8
95% C.I.	( 19.0, 22.1)	( 24.0, 29.0)	( 13.1, 16.7)	( 21.5, 30.7)	( 18.0, 21.2)

	HISP./MULTIRACE NON-WHITE MALES	NON-HISPANIC WHITE MALES	HISP./MULTIRACE NON-WHITE FEMALES	NON-HISPANIC WHITE FEMALES
SAMPLE SIZE	264	1464	310	2104
PREVALENCE	34.6%	24.9%	15.9%	14.8%
STANDARD ERROR	3.7	1.3	2.3	1.0
95% C.I.	( 27.4, 41.8)	( 22.3, 27.5)	( 11.5, 20.3)	( 12.9, 16.7)

	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
<b>MALES/FEMALES</b> *****						
TOTAL	11.6%	1.5	( 8.6,14.6)	7.49*	1.78	( 1.53, 2.07)**
HISP./MULTIRACE NON-WH	18.7%	4.3	(10.2,27.1)	4.33*	2.17	( 1.53, 3.09)**
NON-HISPANIC WHITES	10.0%	1.7	( 6.8,13.3)	6.09*	1.68	( 1.42, 1.98)**
<b>HISP./MULTIRACE NON-WHITES/NON-HISPANIC WHITES</b> *****						
TOTAL	6.5%	2.5	( 1.6,11.4)	2.60*	1.33	( 1.07, 1.65)**
MALES	9.7%	3.9	( 2.1,17.4)	2.49*	1.39	( 1.07, 1.80)**
FEMALES	1.1%	2.5	( 0.0, 5.9)	0.45	1.08	( 0.79, 1.47)

\*DIFFERENCE IS SIGNIFICANT AT ALPHA = .05

\*\*1.0 IS NOT IN THE CONFIDENCE INTERVAL

2001 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM  
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : COLORECTAL CANCER: NO FOBT IN THE PAST TWO YEARS, AGE 50 & OVER (APPROX HP 2010 OBJ. 3.12a)

	TOTAL *****	MALES *****	FEMALES *****	HISP./MULTIRACE NON-WHITE *****	NON-HISPANIC WHITE *****
SAMPLE SIZE	1531	610	921	131	1384
PREVALENCE	59.0%	63.0%	55.6%	59.9%	58.7%
STANDARD ERROR	1.4	2.1	1.8	4.8	1.4
95% C.I.	( 56.4, 61.7)	( 58.9, 67.1)	( 52.1, 59.1)	( 50.6, 69.2)	( 55.9, 61.5)

	HISP./MULTIRACE NON-WHITE MALES	NON-HISPANIC WHITE MALES	HISP./MULTIRACE NON-WHITE FEMALES	NON-HISPANIC WHITE FEMALES
SAMPLE SIZE	63	540	68	844
PREVALENCE	64.8%	62.6%	54.1%	55.5%
STANDARD ERROR	6.5	2.2	6.8	1.9
95% C.I.	( 52.0, 77.6)	( 58.2, 66.9)	( 40.9, 67.4)	( 51.9, 59.1)

	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
<b>MALES/FEMALES</b> *****						
TOTAL	7.4%	2.7	( 2.1,12.8)	2.71*	1.13	( 1.04, 1.24)**
HISP./MULTIRACE NON-WH	10.7%	9.4	( 0.0,29.1)	1.13	1.20	( 0.88, 1.63)
NON-HISPANIC WHITES	7.0%	2.9	( 1.4,12.7)	2.43*	1.13	( 1.02, 1.24)**
<b>HISP./MULTIRACE NON-WHITES/NON-HISPANIC WHITES</b> *****						
TOTAL	1.2%	5.0	( 0.0,10.9)	0.24	1.02	( 0.87, 1.20)
MALES	2.2%	6.9	( 0.0,15.8)	0.32	1.04	( 0.84, 1.28)
FEMALES	1.4%	7.0	( 0.0,15.1)	0.20	0.97	( 0.76, 1.25)

\*DIFFERENCE IS SIGNIFICANT AT ALPHA = .05

\*\*1.0 IS NOT IN THE CONFIDENCE INTERVAL

2001 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM  
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : SIGMOIDOSCOPY: NEVER HAD SIGMOIDOSCOPY, AGE 50 & OVER (HP 2010 OBJ. 3.12b)

	TOTAL *****	MALES *****	FEMALES *****	HISP./MULTIRACE NON-WHITE *****	NON-HISPANIC WHITE *****
SAMPLE SIZE	1533	611	922	128	1389
PREVALENCE	48.0%	48.0%	47.9%	49.4%	47.6%
STANDARD ERROR	1.4	2.2	1.8	4.9	1.5
95% C.I.	( 45.2, 50.7)	( 43.7, 52.3)	( 44.4, 51.4)	( 39.8, 59.0)	( 44.7, 50.5)

  

	HISP./MULTIRACE NON-WHITE MALES	NON-HISPANIC WHITE MALES	HISP./MULTIRACE NON-WHITE FEMALES	NON-HISPANIC WHITE FEMALES
SAMPLE SIZE	61	543	67	846
PREVALENCE	48.2%	47.7%	50.7%	47.5%
STANDARD ERROR	7.0	2.3	6.8	1.9
95% C.I.	( 34.5, 61.9)	( 43.2, 52.2)	( 37.5, 64.0)	( 43.9, 51.2)

	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
<b>MALES/FEMALES</b> *****						
TOTAL	0.1%	2.8	( 0.0, 5.6)	0.04	1.00	( 0.89, 1.12)
HISP./MULTIRACE NON-WH	2.5%	9.7	( 0.0, 21.6)	0.26	0.95	( 0.65, 1.40)
NON-HISPANIC WHITES	0.2%	3.0	( 0.0, 6.0)	0.05	1.00	( 0.89, 1.13)
<b>HISP./MULTIRACE NON-WHITES/NON-HISPANIC WHITES</b> *****						
TOTAL	1.8%	5.1	( 0.0, 11.8)	0.34	1.04	( 0.84, 1.27)
MALES	0.5%	7.4	( 0.0, 14.9)	0.07	1.01	( 0.75, 1.37)
FEMALES	3.2%	7.0	( 0.0, 17.0)	0.46	1.07	( 0.81, 1.41)

\*DIFFERENCE IS SIGNIFICANT AT ALPHA = .05  
\*\*1.0 IS NOT IN THE CONFIDENCE INTERVAL

2001 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM  
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : DIABETES: EVER TOLD YOU HAVE DIABETES BY A DOCTOR, AGE 18 AND OVER (APPROX HP 2010 OBJ. 5.4)

	TOTAL *****	MALES *****	FEMALES *****	HISP./MULTIRACE NON-WHITE *****	NON-HISPANIC WHITE *****
SAMPLE SIZE	4205	1756	2449	575	3578
PREVALENCE	5.7%	6.2%	5.2%	8.4%	5.3%
STANDARD ERROR	0.4	0.6	0.5	1.3	0.4
95% C.I.	( 4.9, 6.5)	( 5.0, 7.4)	( 4.3, 6.2)	( 5.8, 10.9)	( 4.5, 6.0)

  

	HISP./MULTIRACE NON-WHITE MALES	NON-HISPANIC WHITE MALES	HISP./MULTIRACE NON-WHITE FEMALES	NON-HISPANIC WHITE FEMALES
SAMPLE SIZE	265	1468	310	2110
PREVALENCE	8.4%	5.7%	8.3%	4.8%
STANDARD ERROR	1.9	0.6	1.7	0.5
95% C.I.	( 4.7, 12.1)	( 4.5, 7.0)	( 5.0, 11.6)	( 3.8, 5.8)

	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
<b>MALES/FEMALES</b> *****						
TOTAL	0.9%	0.8	( 0.0, 2.5)	1.20	1.18	( 0.90, 1.54)
HISP./MULTIRACE NON-WH	0.1%	2.5	( 0.0, 5.1)	0.05	1.02	( 0.56, 1.84)
NON-HISPANIC WHITES	0.9%	0.8	( 0.0, 2.5)	1.13	1.19	( 0.88, 1.61)
<b>HISP./MULTIRACE NON-WHITES/NON-HISPANIC WHITES</b> *****						
TOTAL	3.1%	1.3	( 0.5, 5.7)	2.31*	1.59	( 1.07, 2.36)**
MALES	2.7%	2.0	( 0.0, 6.6)	1.35	1.47	( 0.84, 2.56)
FEMALES	3.5%	1.8	( 0.0, 7.0)	1.96	1.72	( 1.00, 2.97)

\*DIFFERENCE IS SIGNIFICANT AT ALPHA = .05  
\*\*1.0 IS NOT IN THE CONFIDENCE INTERVAL

2001 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM  
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : CHOLESTEROL SCREENING: NO CHOLESTEROL CHECK IN PAST FIVE YEARS, AGE 18 & OVER (HP 2010 OBJ. 12.15)

	TOTAL *****	MALES *****	FEMALES *****	HISP./MULTIRACE NON-WHITE *****	NON-HISPANIC WHITE *****
SAMPLE SIZE	4058	1700	2358	557	3452
PREVALENCE	28.5%	31.9%	25.4%	35.9%	27.4%
STANDARD ERROR	0.8	1.3	1.1	2.3	0.9
95% C.I.	( 26.9, 30.2)	( 29.3, 34.4)	( 23.3, 27.5)	( 31.3, 40.5)	( 25.6, 29.2)

  

	HISP./MULTIRACE NON-WHITE MALES	NON-HISPANIC WHITE MALES	HISP./MULTIRACE NON-WHITE FEMALES	NON-HISPANIC WHITE FEMALES
SAMPLE SIZE	257	1422	300	2030
PREVALENCE	40.3%	30.4%	30.7%	24.6%
STANDARD ERROR	3.4	1.4	3.1	1.2
95% C.I.	( 33.7, 47.0)	( 27.7, 33.2)	( 24.6, 36.7)	( 22.4, 26.9)

	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
<b>MALES/FEMALES</b> *****						
TOTAL	6.5%	1.7	( 3.2, 9.8)	3.86*	1.26	( 1.12, 1.41)**
HISP./MULTIRACE NON-WH	9.7%	4.6	( 0.6,18.7)	2.10*	1.31	( 1.02, 1.70)**
NON-HISPANIC WHITES	5.8%	1.8	( 2.3, 9.4)	3.20*	1.24	( 1.09, 1.41)**
<b>HISP./MULTIRACE NON-WHITES/NON-HISPANIC WHITES</b> *****						
TOTAL	8.5%	2.5	( 3.6,13.4)	3.39*	1.31	( 1.12, 1.53)**
MALES	9.9%	3.7	( 2.7,17.1)	2.68*	1.32	( 1.08, 1.63)**
FEMALES	6.1%	3.3	( 0.0,12.5)	1.84	1.25	( 0.99, 1.58)

\*DIFFERENCE IS SIGNIFICANT AT ALPHA = .05  
\*\*1.0 IS NOT IN THE CONFIDENCE INTERVAL

2001 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM  
WASHINGTON STATEWIDE

9

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : FLU SHOT: NO FLU SHOT IN PAST YEAR, AGE 65 AND OVER (HP 2010 OBJ. 14.29a)

	TOTAL *****	MALES *****	FEMALES *****	HISP./MULTIRACE NON-WHITE *****	NON-HISPANIC WHITE *****
SAMPLE SIZE	706	256	450	53	649
PREVALENCE	27.5%	28.9%	26.5%	35.4%	26.8%
STANDARD ERROR	1.8	3.0	2.2	7.0	1.9
95% C.I.	( 24.0, 31.1)	( 23.0, 34.8)	( 22.2, 30.8)	( 21.7, 49.1)	( 23.1, 30.5)

	HISP./MULTIRACE NON-WHITE MALES	NON-HISPANIC WHITE MALES	HISP./MULTIRACE NON-WHITE FEMALES	NON-HISPANIC WHITE FEMALES
SAMPLE SIZE	23	230	30	419
PREVALENCE	31.5%	28.6%	39.0%	25.5%
STANDARD ERROR	10.1	3.2	9.6	2.3
95% C.I.	( 11.7, 51.2)	( 22.4, 34.8)	( 20.2, 57.8)	( 21.1, 30.0)

	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
MALES/FEMALES *****						
TOTAL	2.4%	3.7	( 0.0, 9.7)	0.64	1.09	( 0.84, 1.42)
HISP./MULTIRACE NON-WH	7.5%	13.9	( 0.0,34.8)	0.54	0.81	( 0.37, 1.75)
NON-HISPANIC WHITES	3.0%	3.9	( 0.0,10.7)	0.78	1.12	( 0.84, 1.48)
HISP./MULTIRACE NON-WHITES/NON-HISPANIC WHITES *****						
TOTAL	8.6%	7.2	( 0.0,22.8)	1.19	1.32	( 0.83, 2.09)
MALES	2.9%	10.6	( 0.0,23.6)	0.27	1.10	( 0.55, 2.20)
FEMALES	13.5%	9.9	( 0.0,32.8)	1.37	1.53	( 0.83, 2.80)

\*DIFFERENCE IS SIGNIFICANT AT ALPHA = .05  
\*\*1.0 IS NOT IN THE CONFIDENCE INTERVAL

June 18, 2002

2001 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM  
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : PNEUMONIA VACCINATION: NO PNEUMOCOCCAL VACCINATION, AGE 65 AND OVER (HP 2010 OBJ. 14.29b)

	TOTAL *****	MALES *****	FEMALES *****	HISP./MULTIRACE NON-WHITE *****	NON-HISPANIC WHITE *****
SAMPLE SIZE	680	243	437	53	624
PREVALENCE	33.2%	37.5%	30.2%	38.6%	32.7%
STANDARD ERROR	2.0	3.3	2.4	7.1	2.0
95% C.I.	( 29.4, 37.0)	( 31.0, 43.9)	( 25.6, 34.8)	( 24.6, 52.5)	( 28.7, 36.7)

	HISP./MULTIRACE NON-WHITE MALES	NON-HISPANIC WHITE MALES	HISP./MULTIRACE NON-WHITE FEMALES	NON-HISPANIC WHITE FEMALES
SAMPLE SIZE	23	218	30	406
PREVALENCE	31.5%	38.0%	45.0%	29.0%
STANDARD ERROR	10.1	3.5	9.8	2.4
95% C.I.	( 11.7, 51.2)	( 31.1, 44.8)	( 25.8, 64.2)	( 24.3, 33.7)

	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
<b>MALES/FEMALES</b> *****						
TOTAL	7.3%	4.1	( 0.0,15.2)	1.80	1.24	( 0.98, 1.57)
HISP./MULTIRACE NON-WH	13.5%	14.0	( 0.0,41.1)	0.96	0.70	( 0.34, 1.45)
NON-HISPANIC WHITES	9.0%	4.2	( 0.6,17.3)	2.11*	1.31	( 1.02, 1.68)**
<b>HISP./MULTIRACE NON-WHITES/NON-HISPANIC WHITES</b> *****						
TOTAL	5.9%	7.4	( 0.0,20.4)	0.79	1.18	( 0.78, 1.77)
MALES	6.5%	10.7	( 0.0,27.4)	0.61	0.83	( 0.45, 1.52)
FEMALES	16.0%	10.1	( 0.0,35.7)	1.59	1.55	( 0.90, 2.67)

\*DIFFERENCE IS SIGNIFICANT AT ALPHA = .05  
\*\*1.0 IS NOT IN THE CONFIDENCE INTERVAL

2001 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM  
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : OBESITY: BASED ON NEW BODY MASS INDEX (BMI) GREATER THAN OR EQUAL TO 30.0 (APPROX HP 2010 OBJ. 19.2)

	TOTAL *****	MALES *****	FEMALES *****	HISP./MULTIRACE NON-WHITE *****	NON-HISPANIC WHITE *****
SAMPLE SIZE	4026	1734	2292	548	3432
PREVALENCE	19.3%	20.9%	17.8%	21.7%	19.0%
STANDARD ERROR	0.7	1.1	0.9	2.0	0.7
95% C.I.	( 18.0, 20.7)	( 18.8, 22.9)	( 16.1, 19.5)	( 17.8, 25.5)	( 17.6, 20.5)

  

	HISP./MULTIRACE NON-WHITE MALES	NON-HISPANIC WHITE MALES	HISP./MULTIRACE NON-WHITE FEMALES	NON-HISPANIC WHITE FEMALES
SAMPLE SIZE	260	1453	288	1979
PREVALENCE	22.9%	20.5%	20.1%	17.5%
STANDARD ERROR	2.9	1.2	2.6	0.9
95% C.I.	( 17.2, 28.6)	( 18.3, 22.8)	( 15.1, 25.1)	( 15.7, 19.3)

	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
<b>MALES/FEMALES</b> *****						
TOTAL	3.0%	1.4	( 0.4, 5.7)	2.22*	1.17	( 1.02, 1.35)**
HISP./MULTIRACE NON-WH	2.8%	3.9	( 0.0,10.4)	0.71	1.14	( 0.80, 1.62)
NON-HISPANIC WHITES	3.0%	1.5	( 0.1, 5.9)	2.03*	1.17	( 1.01, 1.36)**
<b>HISP./MULTIRACE NON-WHITES/NON-HISPANIC WHITES</b> *****						
TOTAL	2.6%	2.1	( 0.0, 6.8)	1.25	1.14	( 0.93, 1.40)
MALES	2.3%	3.1	( 0.0, 8.5)	0.75	1.11	( 0.84, 1.48)
FEMALES	2.6%	2.7	( 0.0, 7.9)	0.95	1.15	( 0.86, 1.52)

\*DIFFERENCE IS SIGNIFICANT AT ALPHA = .05  
\*\*1.0 IS NOT IN THE CONFIDENCE INTERVAL

2001 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM  
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : LEISURE TIME: NO PHYSICAL ACTIVITY OR EXERCISE DURING PAST THIRTY DAYS, AGE 18 & OVER (HP 2010 OBJ. 22.1)

	TOTAL *****	MALES *****	FEMALES *****	HISP./MULTIRACE NON-WHITE *****	NON-HISPANIC WHITE *****
SAMPLE SIZE	4205	1757	2448	576	3577
PREVALENCE	17.1%	14.9%	19.2%	22.5%	16.1%
STANDARD ERROR	0.6	0.9	0.9	2.0	0.7
95% C.I.	( 15.8, 18.3)	( 13.1, 16.7)	( 17.5, 20.9)	( 18.5, 26.4)	( 14.8, 17.4)

  

	HISP./MULTIRACE NON-WHITE MALES	NON-HISPANIC WHITE MALES	HISP./MULTIRACE NON-WHITE FEMALES	NON-HISPANIC WHITE FEMALES
SAMPLE SIZE	266	1468	310	2109
PREVALENCE	21.1%	13.5%	24.1%	18.4%
STANDARD ERROR	3.0	0.9	2.7	0.9
95% C.I.	( 15.3, 26.8)	( 11.6, 15.4)	( 18.8, 29.5)	( 16.6, 20.3)

	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
<b>MALES/FEMALES</b> *****						
TOTAL	4.3%	1.3	( 1.8, 6.9)	3.37*	0.77	( 0.67, 0.90)**
HISP./MULTIRACE NON-WH	3.1%	4.0	( 0.0,11.0)	0.77	0.87	( 0.62, 1.24)
NON-HISPANIC WHITES	4.9%	1.3	( 2.3, 7.6)	3.69*	0.73	( 0.62, 0.86)**
<b>HISP./MULTIRACE NON-WHITES/NON-HISPANIC WHITES</b> *****						
TOTAL	6.4%	2.1	( 2.2,10.6)	2.98*	1.40	( 1.12, 1.74)**
MALES	7.5%	3.1	( 1.5,13.6)	2.43*	1.56	( 1.09, 2.23)**
FEMALES	5.7%	2.9	( 0.1,11.4)	1.98*	1.31	( 1.00, 1.71)**

\*DIFFERENCE IS SIGNIFICANT AT ALPHA = .05  
\*\*1.0 IS NOT IN THE CONFIDENCE INTERVAL

2001 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM  
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : SMOKING: CURRENT SMOKING STATUS, AGE 18 AND OVER (HP 2010 OBJ. 27.1a)

	TOTAL *****	MALES *****	FEMALES *****	HISP./MULTIRACE NON-WHITE *****	NON-HISPANIC WHITE *****
SAMPLE SIZE	4191	1752	2439	574	3568
PREVALENCE	22.5%	24.5%	20.5%	29.3%	21.2%
STANDARD ERROR	0.7	1.1	0.9	2.2	0.8
95% C.I.	( 21.0, 23.9)	( 22.3, 26.7)	( 18.7, 22.3)	( 25.0, 33.7)	( 19.7, 22.7)

  

	HISP./MULTIRACE NON-WHITE MALES	NON-HISPANIC WHITE MALES	HISP./MULTIRACE NON-WHITE FEMALES	NON-HISPANIC WHITE FEMALES
SAMPLE SIZE	265	1465	309	2103
PREVALENCE	36.1%	22.2%	21.2%	20.4%
STANDARD ERROR	3.4	1.2	2.5	1.0
95% C.I.	( 29.4, 42.8)	( 19.9, 24.5)	( 16.2, 26.2)	( 18.4, 22.3)

	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
<b>MALES/FEMALES</b> *****						
TOTAL	4.0%	1.5	( 1.1, 6.9)	2.74*	1.20	( 1.05, 1.36)**
HISP./MULTIRACE NON-WH	14.9%	4.3	( 6.6, 23.3)	3.50*	1.71	( 1.27, 2.30)**
NON-HISPANIC WHITES	1.9%	1.5	( 0.0, 4.9)	1.21	1.09	( 0.95, 1.26)
<b>HISP./MULTIRACE NON-WHITES/NON-HISPANIC WHITES</b> *****						
TOTAL	8.1%	2.4	( 3.5, 12.7)	3.45*	1.38	( 1.15, 1.66)**
MALES	13.9%	3.6	( 6.8, 21.0)	3.84*	1.63	( 1.27, 2.08)**
FEMALES	0.8%	2.7	( 0.0, 6.2)	0.30	1.04	( 0.80, 1.35)

\*DIFFERENCE IS SIGNIFICANT AT ALPHA = .05  
\*\*1.0 IS NOT IN THE CONFIDENCE INTERVAL

2001 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM  
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : BINGE DRINKING: BINGE DRINKING DURING PAST 30 DAYS, AGE 18 AND OVER (HP 2010 OBJ. 26.11c)

	TOTAL *****	MALES *****	FEMALES *****	HISP./MULTIRACE NON-WHITE *****	NON-HISPANIC WHITE *****
SAMPLE SIZE	4187	1747	2440	575	3560
PREVALENCE	14.9%	23.2%	7.0%	19.3%	14.2%
STANDARD ERROR	0.6	1.1	0.6	2.0	0.7
95% C.I.	( 13.6, 16.1)	( 21.0, 25.4)	( 5.8, 8.1)	( 15.4, 23.2)	( 12.9, 15.6)

	HISP./MULTIRACE NON-WHITE MALES	NON-HISPANIC WHITE MALES	HISP./MULTIRACE NON-WHITE FEMALES	NON-HISPANIC WHITE FEMALES
SAMPLE SIZE	265	1459	310	2101
PREVALENCE	29.3%	22.2%	7.5%	6.9%
STANDARD ERROR	3.2	1.2	1.6	0.6
95% C.I.	( 23.0, 35.6)	( 19.9, 24.5)	( 4.4, 10.7)	( 5.6, 8.2)

	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
<b>MALES/FEMALES</b> *****						
TOTAL	16.2%	1.3	(13.8,18.7)	12.86*	3.33	( 2.77, 4.00)**
HISP./MULTIRACE NON-WH	21.8%	3.6	(14.7,28.8)	6.05*	3.90	( 2.51, 6.06)**
NON-HISPANIC WHITES	15.3%	1.3	(12.7,17.9)	11.37*	3.21	( 2.63, 3.92)**
<b>HISP./MULTIRACE NON-WHITES/NON-HISPANIC WHITES</b> *****						
TOTAL	5.1%	2.1	( 0.9, 9.2)	2.40*	1.35	( 1.06, 1.74)**
MALES	7.1%	3.4	( 0.3,13.8)	2.06*	1.32	( 1.01, 1.71)**
FEMALES	0.6%	1.7	( 0.0, 4.0)	0.34	1.09	( 0.68, 1.74)

\*DIFFERENCE IS SIGNIFICANT AT ALPHA = .05  
\*\*1.0 IS NOT IN THE CONFIDENCE INTERVAL