

Behavioral Risk Factor Surveillance System

State Prevalence Report of Risk Factors for WASHINGTON Statewide 2002



2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
TABLE OF CONTENTS

STATE PREVALENCE REPORT OF RISK FACTORS FOR WASHINGTON STATEWIDE 2002

Table	Page
HEALTH INSURANCE: HAVE NO HEALTH CARE COVERAGE, AGES 18-64.....	3
HEALTH CARE ACCESS: NO SPECIFIC SOURCE OF ON-GOING HEATH CARE, AGES 18 AND OVER.....	4
COLORECTAL SCREENING: NO FOBT IN PAST 2 YRS, AGE 50 AND OVER.....	5
SIGMOIDOSCOPY: NEVER HAD A SIGMOIDOSCOPY/COLONSCOPY, AGE 50 AND OVER.....	6
DIABETES: EVER TOLD YOU HAVE DIABETES BY A DOCTOR.....	7
FLU IMMUNIZATION: NO FLU SHOT IN PAST 12 MONTHS, AGE 65 AND OVER.....	8
PNEUMONIA VACCINATION: DID NOT HAVE A PNEUMONIA VACCINATION, AGE 65 AND OVER.....	9
OBESITY: OBESITY DEFINED BY BODY MASS INDEX, AGE 18 AND OVER.....	10
LEISURE TIME PHYSICAL ACTIVITY: NO PHYSICAL ACTIVITY IN PAST THIRTY DAYS, AGE 18 AND OVER.....	11
TOBACCO USE: CURRENT SMOKER, AGE 18 AND OVER.....	12
ALCOHOL CONSUMPTION: BINGE DRINKING DURING THE PAST THIRTY DAYS, AGE 18 AND OVER.....	13
MAMMOGRAM: DID NOT HAVE A MAMMOGRAM IN PAST TWO YEARS, WOMEN AGE 40 AND OVER.....	14
PAP TEST: NEVER HAD A PAP TEST, WOMEN AGE 18 AND OVER.....	15
PAP SMEAR: DID NOT HAVE A PAP SMEAR WITHIN PAST 3 YEARS, WOMEN AGE 18 AND OVER.....	16

Note :

THIS SECTION OF THE TABULATIONS CONTAINS PREVALENCE ESTIMATES FOR HEALTH BEHAVIORS AND PREVENTIVE HEALTH MEASURES BY RACE AND SEX.

FOR THESE TABLES, THE RACE GROUP 'NON-HISPANIC WHITES' IS DEFINED AS ALL RESPONDENTS WHO INDICATED THEY WERE WHITE AND NOT OF HISPANIC ORIGIN. THE GROUP 'HISPANIC OR NON-WHITES' IS DEFINED AS ALL RACES EXCLUDING NON-HISPANIC WHITE RESPONDENTS.

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR: HEALTH INSURANCE: HAVE NO HEALTH CARE COVERAGE: AGES 18-64 (HP 2010 OBJ. 1.1)

	Total *****	Males *****	Females *****	Hispanic or Multirace Non-White *****	Non-Hispanic White *****
SAMPLE SIZE	3944	1629	2315	634	3277
PREVALENCE	14.0%	16.3%	11.7%	17.8%	13.3%
STANDARD ERROR	0.7	1.1	0.9	1.8	0.8
95% CI	(12.6,15.4)	(14.1,18.6)	(10.0,13.5)	(14.2,21.4)	(11.7,14.8)

	Hispanic or Multirace Non-White Males *****	Non-Hispanic White Males *****	Hispanic or Multirace Non-White Females *****	Non-Hispanic White Females *****
SAMPLE SIZE	284	1330	350	1947
PREVALENCE	22.7%	14.9%	12.1%	11.6%
STANDARD ERROR	3.0	1.2	1.9	1.0
95% CI	(16.9,28.5)	(12.5,17.4)	(8.3,15.8)	(9.6,13.6)

MALES-FEMALE *****	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
TOTAL	4.6%	1.5	(1.8,7.5)	3.18*	1.39	(1.14,1.71)
HISP. NON-WHITE	10.6%	3.5	(3.7,17.5)	3.02*	1.88	(1.25,2.83)**
NON-HISPANIC WHITES	3.3%	1.6	(0.2,6.5)	2.08*	1.29	(1.01,1.63)**

	Non-Hispanic White	Hispanic or Multirace Non-White			
TOTAL	4.6%	2.0	(0.6,8.5)	2.27*	1.35 (1.04,1.74)**
MALES	7.7%	3.2	(1.5,14.0)	2.41*	1.52 (1.08,2.13)**
FEMALES	0.4%	2.2	(0.0,4.7)	0.21	1.04 (0.73,1.48)

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05
** 1.0 IS NOT IN THE CONFIDENCE INTERVAL

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR: NO SPECIFIC SOURCE OF ON-GOING HEALTH CARE: AGE 18 & OVER (HP 2010 OBJ. 1.4c)

	Total *****	Males *****	Females *****	Hispanic or Multirace Non-White *****	Non-Hispanic White *****
SAMPLE SIZE	4873	1958	2915	701	4128
PREVALENCE	19.5%	25.3%	13.8%	22.2%	19.0%
STANDARD ERROR	0.8	1.3	0.8	2.0	0.8
95% CI	(18.0,21.0)	(22.8,27.8)	(12.2,15.4)	(18.3,26.0)	(17.3,20.6)

	Hispanic or Multirace Non-White Males *****	Non-Hispanic White Males *****	Hispanic or Multirace Non-White Females *****	Non-Hispanic White Females *****
SAMPLE SIZE	310	1627	391	2501
PREVALENCE	29.0%	24.7%	14.1%	13.6%
STANDARD ERROR	3.1	1.4	2.0	0.9
95% CI	(22.9,35.2)	(22.0,27.5)	(10.3,17.9)	(11.8,15.4)

MALES-FEMALE *****	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
TOTAL	11.5%	1.5	(8.5,14.5)	7.56*	1.84	(1.57,2.15)
HISP. NON-WHITE	14.9%	3.7	(7.7,22.2)	4.03*	2.06	(1.45,2.92)**
NON-HISPANIC WHITES	11.1%	1.7	(7.8,14.4)	6.58*	1.81	(1.52,2.17)**

	Non-Hispanic White	Hispanic or Multirace Non-White			
TOTAL	3.2%	2.1	(0.0,7.4)	1.50	1.17 (0.95,1.43)
MALES	4.3%	3.4	(0.0,11.0)	1.25	1.17 (0.91,1.51)
FEMALES	0.5%	2.2	(0.0,4.7)	0.22	1.04 (0.76,1.41)

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05
** 1.0 IS NOT IN THE CONFIDENCE INTERVAL

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR: COLORECTAL CANCER: NO FOBT IN THE PAST TWO YEARS, AGE 50 & OVER (APPROX HP 2010 OBJ. 3.12a)

	Total *****	Males *****	Females *****	Hispanic or Multirace Non-White *****	Non-Hispanic White *****
SAMPLE SIZE	2054	795	1259	190	1844
PREVALENCE	63.5%	68.8%	58.9%	70.4%	62.5%
STANDARD ERROR	1.3	1.9	1.7	3.8	1.4
95% CI	(61.0,66.0)	(65.0,72.5)	(55.5,62.2)	(63.0,77.7)	(59.8,65.3)

	Hispanic or Multirace Non-White Males *****	Non-Hispanic White Males *****	Hispanic or Multirace Non-White Females *****	Non-Hispanic White Females *****
SAMPLE SIZE	76	709	114	1135
PREVALENCE	72.2%	68.2%	68.5%	57.7%
STANDARD ERROR	5.7	2.1	4.8	1.8
95% CI	(61.0,83.4)	(64.2,72.2)	(59.1,77.9)	(54.1,61.3)

MALES-FEMALE *****	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
TOTAL	9.9%	2.6	(4.9,14.9)	3.85*	1.17	(1.08,1.26)
HISP. NON-WHITE	3.7%	7.5	(0.0,18.3)	0.50	1.05	(0.86,1.30)
NON-HISPANIC WHITES	10.5%	2.8	(5.1,15.9)	3.81*	1.18	(1.08,1.29)**

	Non-Hispanic White	Hispanic or Multirace Non-White				
TOTAL	7.8%	4.0	(0.0,15.7)	1.95	1.13	(1.00,1.27)
MALES	4.0%	6.1	(0.0,15.9)	0.66	1.06	(0.89,1.26)
FEMALES	10.8%	5.1	(0.8,20.8)	2.11*	1.19	(1.01,1.39)**

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05
** 1.0 IS NOT IN THE CONFIDENCE INTERVAL

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR: SIGMOIDOSCOPY: NEVER HAD SIGMOIDOSCOPY, AGE 50 & OVER (HP 2010 OBJ. 3.12b)

	Total *****	Males *****	Females *****	Hispanic or Multirace Non-White *****	Non-Hispanic White *****
SAMPLE SIZE	2070	803	1267	193	1859
PREVALENCE	45.3%	47.2%	43.5%	51.4%	44.4%
STANDARD ERROR	1.4	2.2	1.7	4.4	1.4
95% CI	(42.6,47.9)	(43.0,51.5)	(40.2,46.9)	(42.7,60.0)	(41.6,47.2)

	Hispanic or Multirace Non-White Males *****	Non-Hispanic White Males *****	Hispanic or Multirace Non-White Females *****	Non-Hispanic White Females *****
SAMPLE SIZE	81	712	112	1147
PREVALENCE	54.7%	45.9%	47.8%	43.1%
STANDARD ERROR	6.9	2.3	5.3	1.8
95% CI	(41.2,68.2)	(41.4,50.4)	(37.4,58.2)	(39.5,46.6)

MALES-FEMALE *****	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
TOTAL	3.7%	2.8	(0.0,9.1)	1.33	1.08	(0.96,1.22)
HISP. NON-WHITE	6.9%	8.7	(0.0,23.9)	0.79	1.14	(0.82,1.60)
NON-HISPANIC WHITES	2.8%	2.9	(0.0,8.6)	0.97	1.07	(0.94,1.21)

	Non-Hispanic White	Hispanic or Multirace Non-White				
TOTAL	7.0%	4.7	(0.0,16.1)	1.49	1.16	(0.96,1.40)
MALES	8.7%	7.3	(0.0,23.0)	1.20	1.19	(0.90,1.58)
FEMALES	4.7%	5.6	(0.0,15.7)	0.83	1.11	(0.87,1.41)

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05
** 1.0 IS NOT IN THE CONFIDENCE INTERVAL

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR: DIABETES: EVER TOLD YOU HAVE DIABETES BY A DOCTOR, AGE 18 AND OVER (APPROX HP 2010 OBJ. 5.4)

	Total *****	Males *****	Females *****	Hispanic or Multirace Non-White *****	Non-Hispanic White *****
SAMPLE SIZE	4882	1963	2919	703	4133
PREVALENCE	5.8%	6.4%	5.2%	5.9%	5.7%
STANDARD ERROR	0.4	0.7	0.5	0.9	0.5
95% CI	(5.0,6.6)	(5.0,7.7)	(4.3,6.1)	(4.0,7.7)	(4.8,6.6)

	Hispanic or Multirace Non-White Males *****	Non-Hispanic White Males *****	Hispanic or Multirace Non-White Females *****	Non-Hispanic White Females *****
SAMPLE SIZE	312	1628	391	2505
PREVALENCE	4.5%	6.7%	7.5%	4.8%
STANDARD ERROR	1.1	0.8	1.5	0.5
95% CI	(2.3,6.8)	(5.1,8.2)	(4.4,10.5)	(3.9,5.7)

MALES-FEMALE *****	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
TOTAL	1.2%	0.8	(0.0,2.8)	1.43	1.23	(0.93,1.62)
HISP. NON-WHITE	2.9%	1.9	(0.0,6.7)	1.54	0.61	(0.32,1.15)
NON-HISPANIC WHITES	1.9%	0.9	(0.1,3.7)	2.07*	1.40	(1.02,1.93)**

	Non-Hispanic White	Hispanic or Multirace Non-White			
TOTAL	0.2%	1.0	(0.0,2.2)	0.17	1.03 (0.72,1.47)
MALES	2.2%	1.4	(0.0,4.9)	1.56	0.68 (0.41,1.11)
FEMALES	2.7%	1.6	(0.0,5.8)	1.67	1.56 (0.93,2.64)

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05

** 1.0 IS NOT IN THE CONFIDENCE INTERVAL

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR: FLU SHOT: NO FLU SHOT IN PAST YEAR, AGE 65 AND OVER (HP 2010 OBJ. 14.29a)

	Total *****	Males *****	Females *****	Hispanic or Multirace Non-White *****	Non-Hispanic White *****
SAMPLE SIZE	918	324	594	66	842
PREVALENCE	35.0%	40.6%	30.8%	35.5%	34.1%
STANDARD ERROR	1.9	3.2	2.3	6.9	2.0
95% CI	(31.2,38.7)	(34.3,46.9)	(26.4,35.3)	(22.0,49.0)	(30.2,38.0)

	Hispanic or Multirace Non-White Males *****	Non-Hispanic White Males *****	Hispanic or Multirace Non-White Females *****	Non-Hispanic White Females *****
SAMPLE SIZE	26	292	40	550
PREVALENCE	38.3%	39.2%	32.7%	30.4%
STANDARD ERROR	11.1	3.4	8.1	2.3
95% CI	(16.6,60.0)	(32.6,45.9)	(16.9,48.5)	(25.9,35.0)

MALES-FEMALE *****	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
TOTAL	9.7%	3.9	(2.0,17.5)	2.47*	1.32	(1.06,1.64)
HISP. NON-WHITE	5.6%	13.7	(0.0,32.5)	0.41	1.17	(0.55,2.50)
NON-HISPANIC WHITES	8.8%	4.1	(0.7,16.9)	2.14*	1.29	(1.02,1.63)**

	Non-Hispanic White	Hispanic or Multirace Non-White				
TOTAL	1.4%	7.2	(0.0,15.5)	0.20	1.04	(0.70,1.56)
MALES	0.9%	11.6	(0.0,23.7)	0.08	0.98	(0.54,1.75)
FEMALES	2.3%	8.4	(0.0,18.7)	0.27	1.07	(0.64,1.81)

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05
** 1.0 IS NOT IN THE CONFIDENCE INTERVAL

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR: PNEUMONIA VACCINATION: NO PNEUMOCOCCAL VACCINATION, AGE 65 AND OVER (HP 2010 OBJ. 14.29b)

	Total *****	Males *****	Females *****	Hispanic or Multirace Non-White *****	Non-Hispanic White *****
SAMPLE SIZE	889	304	585	62	817
PREVALENCE	37.0%	42.0%	33.5%	34.6%	36.8%
STANDARD ERROR	2.0	3.3	2.5	6.5	2.1
95% CI	(33.1,40.9)	(35.6,48.4)	(28.6,38.5)	(21.8,47.5)	(32.7,40.9)

	Hispanic or Multirace Non-White Males *****	Non-Hispanic White Males *****	Hispanic or Multirace Non-White Females *****	Non-Hispanic White Females *****
SAMPLE SIZE	24	274	38	543
PREVALENCE	35.9%	41.3%	33.4%	33.8%
STANDARD ERROR	10.3	3.4	8.2	2.6
95% CI	(15.7,56.2)	(34.6,48.0)	(17.3,49.4)	(28.7,39.0)

MALES-FEMALE *****	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
TOTAL	8.4%	4.1	(0.4,16.5)	2.05*	1.25	(1.01,1.55)
HISP. NON-WHITE	2.6%	13.2	(0.0,28.4)	0.19	1.08	(0.51,2.27)
NON-HISPANIC WHITES	7.5%	4.3	(0.0,16.0)	1.73	1.22	(0.97,1.53)

	Non-Hispanic White	Hispanic or Multirace Non-White			
TOTAL	2.2%	6.9	(0.0,15.6)	0.32	0.94 (0.65,1.37)
MALES	5.4%	10.9	(0.0,26.7)	0.49	0.87 (0.50,1.51)
FEMALES	0.5%	8.6	(0.0,17.3)	0.05	0.99 (0.60,1.63)

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05
** 1.0 IS NOT IN THE CONFIDENCE INTERVAL

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR: OBESITY: BASED ON NEW BODY MASS INDEX (BMI) GREATER THAN OR EQUAL TO 30.0 (APPROX HP 2010 OBJ. 19.2)

	Total *****	Males *****	Females *****	Hispanic or Multirace Non-White *****	Non-Hispanic White *****
SAMPLE SIZE	4674	1942	2732	667	3962
PREVALENCE	21.3%	21.2%	21.3%	23.4%	20.9%
STANDARD ERROR	0.8	1.2	0.9	2.0	0.8
95% CI	(19.8,22.7)	(18.9,23.5)	(19.5,23.2)	(19.4,27.4)	(19.3,22.5)

	Hispanic or Multirace Non-White Males *****	Non-Hispanic White Males *****	Hispanic or Multirace Non-White Females *****	Non-Hispanic White Females *****
SAMPLE SIZE	303	1616	364	2346
PREVALENCE	21.0%	21.2%	26.2%	20.6%
STANDARD ERROR	2.9	1.3	2.7	1.0
95% CI	(15.3,26.8)	(18.6,23.7)	(20.9,31.6)	(18.6,22.5)

MALES-FEMALE *****	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
TOTAL	0.1%	1.5	(0.0,3.1)	0.09	0.99	(0.86,1.14)
HISP. NON-WHITE	5.2%	4.0	(0.0,13.1)	1.30	0.80	(0.57,1.12)
NON-HISPANIC WHITES	0.6%	1.6	(0.0,3.8)	0.37	1.03	(0.88,1.20)

	Non-Hispanic White	Hispanic or Multirace Non-White				
TOTAL	2.5%	2.2	(0.0,6.8)	1.16	1.12	(0.92,1.36)
MALES	0.1%	3.2	(0.0,6.4)	0.04	0.99	(0.74,1.34)
FEMALES	5.7%	2.9	(0.0,11.4)	1.96*	1.28	(1.00,1.63)**

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05
** 1.0 IS NOT IN THE CONFIDENCE INTERVAL

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR: LEISURE TIME: NO PHYSICAL ACTIVITY OR EXERCISE DURING PAST THIRTY DAYS, AGE 18 & OVER (HP 2010 OBJ. 22.1)

	Total *****	Males *****	Females *****	Hispanic or Multirace Non-White *****	Non-Hispanic White *****
SAMPLE SIZE	4881	1965	2916	701	4134
PREVALENCE	15.0%	13.6%	16.3%	21.9%	13.7%
STANDARD ERROR	0.7	1.0	0.9	2.0	0.7
95% CI	(13.7,16.3)	(11.6,15.6)	(14.7,18.0)	(18.0,25.8)	(12.4,15.1)

	Hispanic or Multirace Non-White Males *****	Non-Hispanic White Males *****	Hispanic or Multirace Non-White Females *****	Non-Hispanic White Females *****
SAMPLE SIZE	310	1632	391	2502
PREVALENCE	15.0%	13.4%	30.1%	14.0%
STANDARD ERROR	2.5	1.1	3.1	0.8
95% CI	(10.2,19.8)	(11.2,15.6)	(24.0,36.1)	(12.4,15.6)

MALES-FEMALE *****	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
TOTAL	2.7%	1.3	(0.1,5.3)	2.07*	0.83	(0.70,0.99)
HISP. NON-WHITE	15.1%	3.9	(7.4,22.8)	3.83*	0.50	(0.35,0.71)**
NON-HISPANIC WHITES	0.6%	1.4	(0.0,3.4)	0.46	0.95	(0.78,1.16)

	Non-Hispanic White	Hispanic or Multirace Non-White				

TOTAL	8.2%	2.1	(4.1,12.4)	3.90*	1.60	(1.26,2.03)**
MALES	1.6%	2.7	(0.0,6.9)	0.59	1.12	(0.77,1.62)
FEMALES	16.1%	3.2	(9.8,22.3)	5.03*	2.15	(1.59,2.89)**

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05
** 1.0 IS NOT IN THE CONFIDENCE INTERVAL

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR: SMOKING: CURRENT SMOKING STATUS, AGE 18 AND OVER (HP 2010 OBJ. 27.1a)

	Total *****	Males *****	Females *****	Hispanic or Multirace Non-White *****	Non-Hispanic White *****
SAMPLE SIZE	4863	1959	2904	700	4119
PREVALENCE	21.5%	23.6%	19.4%	21.8%	21.5%
STANDARD ERROR	0.8	1.2	0.9	1.9	0.9
95% CI	(19.9,23.0)	(21.2,26.0)	(17.6,21.2)	(18.0,25.6)	(19.8,23.2)

	Hispanic or Multirace Non-White Males *****	Non-Hispanic White Males *****	Hispanic or Multirace Non-White Females *****	Non-Hispanic White Females *****
SAMPLE SIZE	311	1626	389	2493
PREVALENCE	22.3%	24.0%	21.2%	19.2%
STANDARD ERROR	2.8	1.4	2.6	1.0
95% CI	(16.8,27.7)	(21.3,26.8)	(16.0,26.3)	(17.2,21.1)

MALES-FEMALE *****	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
TOTAL	4.2%	1.6	(1.2,7.2)	2.70*	1.22	(1.06,1.40)
HISP. NON-WHITE	1.1%	3.8	(0.0,8.7)	0.30	1.05	(0.75,1.49)
NON-HISPANIC WHITES	4.9%	1.7	(1.5,8.2)	2.83*	1.25	(1.07,1.47)**

	Non-Hispanic White	Hispanic or Multirace Non-White			
TOTAL	0.3%	2.1	(0.0,4.4)	0.13	1.01 (0.84,1.23)
MALES	1.7%	3.1	(0.0,7.8)	0.56	0.93 (0.71,1.21)
FEMALES	2.0%	2.8	(0.0,7.5)	0.71	1.10 (0.84,1.45)

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05
** 1.0 IS NOT IN THE CONFIDENCE INTERVAL

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR: BINGE DRINKING: BINGE DRINKING DURING PAST 30 DAYS, AGE 18 AND OVER (HP 2010 OBJ. 26.11c)

	Total *****	Males *****	Females *****	Hispanic or Multirace Non-White *****	Non-Hispanic White *****
SAMPLE SIZE	4858	1953	2905	699	4115
PREVALENCE	15.1%	22.2%	8.2%	13.9%	15.4%
STANDARD ERROR	0.7	1.1	0.7	1.7	0.7
95% CI	(13.8,16.4)	(19.9,24.4)	(6.9,9.5)	(10.6,17.1)	(14.0,16.8)

	Hispanic or Multirace Non-White Males *****	Non-Hispanic White Males *****	Hispanic or Multirace Non-White Females *****	Non-Hispanic White Females *****
SAMPLE SIZE	309	1623	390	2492
PREVALENCE	20.9%	22.6%	5.7%	8.7%
STANDARD ERROR	2.8	1.3	1.2	0.7
95% CI	(15.4,26.4)	(20.2,25.1)	(3.4,8.1)	(7.3,10.2)

MALES-FEMALE *****	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
TOTAL	13.9%	1.3	(11.4,16.5)	10.67*	2.70	(2.25,3.24)
HISP. NON-WHITE	15.1%	3.1	(9.2,21.1)	4.96*	3.65	(2.19,6.08)**
NON-HISPANIC WHITES	13.9%	1.5	(11.0,16.7)	9.54*	2.59	(2.13,3.15)**

	Non-Hispanic White	Hispanic or Multirace Non-White				
TOTAL	1.5%	1.8	(0.0,5.1)	0.84	0.90	(0.71,1.15)
MALES	1.8%	3.1	(0.0,7.8)	0.57	0.92	(0.70,1.22)
FEMALES	3.0%	1.4	(0.2,5.8)	2.13*	0.66	(0.44,0.97)**

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05
** 1.0 IS NOT IN THE CONFIDENCE INTERVAL

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : MAMMOGRAM: DID NOT HAVE A MAMMOGRAM IN PAST TWO YEARS, WOMEN AGE 40 AND OVER (HP 2010 OBJ. 3.13)

	Total	Hispanic or Multirace Non-White	Non-Hispanic White
*****	*****	*****	*****
SAMPLE SIZE	1890	194	1680
PREVALENCE	25.9%	29.9%	25.3%
STANDARD ERROR	1.2	3.8	1.3
95% CI	(23.4,28.3)	(22.4,37.4)	(22.7,27.9)

Hispanic or Multirace
Non-White

*****	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
Non-Hispanic White *****						
FEMALES	5.0%	4.0	.	1.00	1.00	(0.9,1.6)0

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05

** 1.0 IS NOT IN THE CONFIDENCE INTERVAL

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : PAP SMEAR: NEVER HAD A PAP, WOMEN AGE 18 AND OVER (APPROX HP 2010 OBJ 3.11a)

	Total	Hispanic or Multirace Non-White	Non-Hispanic White
*****	*****	*****	*****
SAMPLE SIZE	2919	391	2505
PREVALENCE	4.2%	6.8%	3.8%
STANDARD ERROR	0.5	2.3	0.5
95% CI	(3.1,5.3)	(2.4,11.2)	(2.8,4.8)

Hispanic or Multirace
Non-White

*****	ABSOLUTE DIFFERENCE *****	STANDARD ERROR FOR DIFFERENCE *****	95% C.I. FOR DIFFERENCE *****	Z-STATISTIC FOR DIFFERENCE *****	PREVALENCE RATIO *****	95% C.I. FOR PREVALENCE RATIO *****
Non-Hispanic White *****						
FEMALES	3.0%	2.0	.	1.00	2.00	(0.7,4.3)0

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05
** 1.0 IS NOT IN THE CONFIDENCE INTERVAL

2002 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM
WASHINGTON STATEWIDE

PREVALENCE REPORT OF RISK FACTORS BY SEX AND RACE

RISK FACTOR : PAP SMEAR: NO PAP SMEAR IN PAST 3 YRS, WOMEN AGE 18 AND OVER (APPROX HP 2010 OBJ 3.11b)

	Total	Hispanic or Multirace Non-White	Non-Hispanic White
*****	*****	*****	*****
SAMPLE SIZE	2919	391	2505
PREVALENCE	17%	17.1%	17.1%
STANDARD ERROR	0.9	2.7	0.9
95% CI	(15.3,18.7)	(11.8,22.5)	(15.3,18.8)

Hispanic or Multirace
Non-White

	ABSOLUTE DIFFERENCE	STANDARD ERROR FOR DIFFERENCE	95% C.I. FOR DIFFERENCE	Z-STATISTIC FOR DIFFERENCE	PREVALENCE RATIO	95% C.I. FOR PREVALENCE RATIO
*****	*****	*****	*****	*****	*****	*****
Non-Hispanic White						
*****	*****	*****	*****	*****	*****	*****
FEMALES	0.0%	3.0	.	0.00	1.00	(0.7,1.4)0

* DIFFERENCE IS SIGNIFICANT AT ALPHA=.05

** 1.0 IS NOT IN THE CONFIDENCE INTERVAL