



Cudurka Balaygga Sambabada (Pneumonic Plague)

Waa maxay cudurka balaygga?

Balaygu waa cudur aan caam ahayn oo la kala qaado oo ku dhaca xayawaanka iyo bini'aadanka oo uu keeno bakteeriyyada loo yaqaano *Yersinia pestis* (*Y. pestis*).

Bakteeriyyada *Y. pestis* waxay ku jirtaa jiirka ama xayawaanka la mid ah ee duurjoogto ah iyo takfigooda ama injirboodadooda kuwaasoo laga helo meelo badan oo adduunka ah, oo ay ku jiraan badi galbeedka Maraykanka.

Noocyada balaygga

Balaygga waa la isu tabin karaa waxana uu keeni karaa cudur ah mid ka mid ah qaababkan ama ka badan:

- Marka bakteeriyyada *Y. pestis* jidhka ka soo gasho barta uu ka qaniinay takfi ama injirboodo cudureysan ama marka uu ka soo galo meel jeexan ama dilaacsan oo ah maqaarka, cudurka qofka ku dhaca waxa loo yaqaana buuboonik balayg (**bubonic plague**). “Buboies” waa nudo limfi (lymph nodes) oo bararsan oo xanuun leh. Buuboonik balayg waa nooca ugu caamsan ama ugu badan cudurka isla markaana haddii aan la daweyn waxa uu isu rogi karaa balaygga dhiig-sumowga (septicemic plague) (hoos ka fiiri).
- Marka bakteeriyyada *Y. pestis* ku ururto dhiigga, waxa dhaca shoog dhiig-sumow (septic shock) waxana cudurka sidaas ku yimaada loo yaqaana balaygga dhiig-sumow (**septicemic plague**).
- Marka bakteeriyyada *Y. pestis* la neefsado, waxay galaan sambabada waxana cudurka sidaas ku yimaada loo yaqaana balaygga sambabada (**pneumonic plague**). Balaygga sambababdu waxa uu leeyahay walaac gaar ah sababta oo ah qofku waxa uu qaadsiin karaa dad kale.

Sidee buu balaygga sambababdu u fidaa?

Waa maxay astaamuhu?

Balaygga sambababdu waxa uu yimaadaa marka bakteeriyyada *Y. pestis* la neefsado. Waxa cudurku ku fidi karaa isu soo dhawaansha ah fool-ka-fool marka qofka cudureysan uu qufaco ama hindhisoo. Sababta oo ah waxa uu jidhka galaa marka la neefsado ama neefta la jiido, balaygga sambabada waxa loo fidin karaa si kas ama ogaan ah haddii bakteeriyyada lagu rido weel buufin.

Astaamaha balaygga sambabada waxay bilaabmaan hal ilaa afar maalmood ka dib soo gaadhista bakteeriyyada. Astaamaha waxa ka mid ah qandho, madax-xanuun, tabar-dari iyo qufac dhiig leh ama biyo-biyo ah sababta oo ah caabuqa ama infekshanka sambabada (pneumonia). Caabuqa sambabada ayaa dhakhso u sii xumaada oo — haddii aan goor hore la daweyn —waxa uu noqon karaa mid keena dhimasho.

Talaabooyin ka-hortagga

Haddii aad leedahay astaamo, sida ugu dhakhsaha badan ula tasho bixiye daryeel caafimaad. Haddii aad rumeysan tahay in si kas ama ogaan ah laguu soo gaadhsiiyay balaygga sambabada, waa inaad isla markiiba la xidhiidhid saraakiisha fulinta sharciga.

Ma jiro tallaal lid ku ah ama wax ka tara balaygga sambabada. Waxa la isticmaalaa antibiyootiko si looga hortago jirro ku dhacda kuwa uu soo gaadhaday balaygga sambababdu.

Daweynta balaygga sambabada

Daweyn goor hore ah oo la isticmaalayo antibiyootiko haboon ayaa muhiim ah sababta oo ah balaygga aan la daweyn—gaar ahaan nooca sambabada—badi markasta waa loo dhintaa. Waa inaad isticmaashid antibiyootiko si looga hortago ama loo daweyno balaygga kaliya iyada oo uu ku maareynayo bixiyahaaga daryeel caafimaad ama waaxda caafimaadka ee degmada.



PO Box 47890
Olympia, WA 98504-
7890

Bogga Internetka:
www.doh.wa.gov

360-236-4027
(800) 525-0127

DOH Pub 821-023
Somali