

# Chronic Disease Profile

## Introduction

This report summarizes key health statistics related to chronic disease burden and risk for local populations in Washington State. The Department of Health uses established population and health surveillance systems to describe the current prevalence<sup>i</sup> of important health indicators within specific populations, and also to provide comparisons of the prevalence within specific populations to the state overall.

These data can be used to plan interventions or describe the importance and need for health interventions. Interventions may be directed to specific health conditions, or to factors that impact many aspects of health, such as income, education and housing. Therefore, this report may be useful for community members, leaders or other stakeholders who are working to improve the health status of the community.

## Life Course Approach<sup>ii</sup>

Health and quality of life at all stages in life depend on the cumulative effects of behaviors and exposures earlier in life, and on social, genetic, and epigenetic effects<sup>iii</sup> that span generations. A mother's experiences, even prior to conception, can alter the development of the fetus and child. Choices made by adolescents grow out of the experiences of childhood, and can shape behavior later in adulthood. A lifetime of risky behavior or exposure to toxic or stressful conditions can lead to chronic disease, poor quality of life and early death.

This report is organized based on a life course approach. We begin with data on the demographic, social and economic context. Next we show data related to birth and early childhood. We follow these in turn by data for youth (grade 10), adults (age 18+) and seniors (age 65+). Lastly, we provide patterns of mortality.

## Health Risk Indicators

Many pieces of health data can be presented in either a positive or negative manner. For example, we could either talk about reducing obesity, or achieving healthy weight. For other data, only the negative presentation makes sense. For example, it would be awkward to discuss increasing the prevalence of people without diabetes. For consistency and ease of comparison, this document presents all data in terms of risk.

Health data are estimated with some degree of statistical uncertainty. We present the degree of uncertainty by surrounding each estimate in graphs with error bars that represent the 95% confidence interval. See appendix for further detail.

Data sources, explanatory notes, and a glossary of terms are provided in the appendix.

## Racial / Ethnic Designations

Race and ethnicity for most indicators are determined based on self-report by survey respondent. For birth and Pregnancy Risk Assessment Monitoring System (PRAMS) data, race and ethnicity are designated based on the mother's reported race/ethnicity. For death data, race/ethnicity is based on the report of next of kin, or in some cases, by the certifying physician. For consistency with other agency reports, Hispanic ethnicity is treated as a separate race. Respondents who identify themselves as both White and Hispanic are classified as Hispanic. Surveys and data collection systems do not all characterize race and ethnicity in the same way. In some surveys multiracial respondents are classified as a separate group; in others multiracial respondents are asked to choose a single preferred race.

- Census Bureau, American Community Survey (ACS): Respondents identify their ethnicity as Hispanic/Latino or not Hispanic/Latino. Respondents then identify their race by choosing any that apply from a detailed list of racial categories. Identification as multiracial is allowed. In this report, non-Hispanic White refers to those who select non-Hispanic ethnicity, and select White as their only racial classification. Multiracial respondents are not included within non-Hispanic White.
- Birth certificates / PRAMS: Race/Hispanic origin for the mother and father are collected by asking the mother for the data. In this report, we report health indicators according to the mother's race. Mothers choose any that apply from a detailed list of racial and ethnic categories. Identification as multiracial is allowed. In this report, non-Hispanic White refers to those who select non-Hispanic ethnicity, and select White as their only racial classification. Multiracial respondents are not included within non-Hispanic White.
- Behavioral Risk Factor Surveillance System (BRFSS): Respondents first identify their ethnicity as Hispanic/Latino or not Hispanic/Latino. Respondents then identify their race by choosing one response from a list of racial categories. Respondents who identify themselves as multiracial are then asked to choose a single preferred racial classification. In this report, non-Hispanic White refers to those who select non-Hispanic ethnicity, and then select White as their preferred race.
- Healthy Youth Survey (HYS): Respondents are asked, "How do you describe yourself? (Select one or more responses.)" Response options are: a. American Indian or Alaskan Native; b. Asian or Asian American; c. Black or African-American; d. Hispanic or Latino/Latina; e. Native Hawaiian or other Pacific Islander; f. White or Caucasian; g. Other. Respondents who check more than one option are classified as multiracial. In this report White refers to those who identify themselves only as White.
- Death certificates: Reporting of race/Hispanic origin on death certificates is sometimes based on observing the decedent, rather than questioning the next of kin. This procedure causes an underestimate of deaths for certain groups, particularly Native Americans, some of the Asian subgroups and Hispanics. Identification as multiracial is allowed. In this report, non-Hispanic White refers to those who are identified as non-Hispanic ethnicity, and as White as their only racial classification. Multiracial decedents are not included within non-Hispanic White.


## Population

**White Population: 5,005,607 = 72% of state**

### Age Distribution


- Washington State: 14% are age 65+; 23% are age < 18
- White: 17% are age 65+; 19% are age < 18

### Population by Race / Ethnicity



**Data Source:** Washington State Office of Financial Management, Forecasting Division, single year intercensal estimates, 2016

## Social and Economic Risk Factors




### Indicator Notes

1. Federal Poverty Level (FPL) is determined based on household income and household size. In 2015, the federal poverty level household income for a family of four was \$24,250.
2. Highest educational attainment is among adults 25 and older.
3. Health Insurance: Did not have any form of health insurance among adults age 18 to 64.

**Data Source:** US Census Bureau, American Community Survey (ACS) Public Use Microdata Sample, 2015

## Prenatal and Birth Health Risk Factors



### Birth Rate

- Washington State: 64 births per 1000 reproductive age women (age 15-44)
- White: 60 births per 1000 reproductive age women (age 15-44)


### Indicator Notes

1. Third trimester smoking: Smoked one or more cigarettes on an average day during the last three months of pregnancy.
2. Prenatal care includes visits to a doctor, nurse, or other healthcare worker before the baby was born to get checkups and advice about pregnancy.
3. Unintended pregnancy: When asked “Thinking back to just before you got pregnant with your new baby, how did you feel about becoming pregnant?” responded “I wanted to become pregnant later” or “I didn’t want to be pregnant then, or in the future.”
4. Low birth weight is defined as a birth under 2,500g but no lighter than 227g. Infants born less than 227g are considered pre-viable.
5. Premature delivery is defined as gestation < 37 weeks.
6. Adolescent mother is defined as age 15-17.

### Data Sources:

- Prenatal: Washington State Department of Health, Pregnancy Risk Assessment Monitoring System (PRAMS), 2012-2014
- Birth outcomes: Washington State Department of Health, Center for Health Statistics (CHS), Birth Certificate Data, 2011-2015.
- Birth rate: Washington State Department of Health, Center for Health Statistics (CHS), Birth Certificate Data, 2015.

## Early Childhood Health Risk Factors




### Indicator Notes

1. Breastfeeding: did not breastfeed baby, or breastfed for less than 8 weeks.
2. Child poverty: Age 0-4, living in a household with income less than FPL.

### Data Sources:

- Child poverty: American Community Survey 2011-2015
- Breastfeeding: Pregnancy Risk Assessment Monitoring System 2012-2014

## Youth (10<sup>th</sup> grade) Health Risk Behaviors




### Indicator Notes

1. Youth smoking, marijuana, alcohol, e-cigarettes: Students are asked “during the past 30 days, how many times did you... Smoke cigarettes; Use marijuana or hashish (grass, hash, pot); Drink a glass, can, or bottle of alcohol (beer, wine, wine coolers, hard liquor); use electronic cigarettes or e-cigs?”
2. The Centers for Disease Control and Prevention (CDC) recommends 60 minutes moderate or vigorous physical activity every day for youths.
3. Poor nutrition is indicated by eating fruits and vegetables less than once a day.

**Data Source:** Washington State Healthy Youth Survey 2016.

## Youth (10<sup>th</sup> grade) Health Risk Conditions




### Indicator Notes

1. Youth obesity: Youth are classified as obese if they are in the 95<sup>th</sup> percentile for body mass index by age and sex based on growth charts developed by the CDC (2000).
2. Bullied: Students are asked “A student is being bullied when another student, or group of students, say or do nasty or unpleasant things to him or her. It is also bullying when a student is teased repeatedly in a way he or she doesn’t like. It is NOT bullying when two students of about the same strength argue or fight. In the last 30 days, how often have you been bullied?”
3. Depression: Students were asked “During the past 12 months, did you ever feel so sad or hopeless almost every day for two weeks or more in a row that you stopped doing some usual activities?”
4. Suicide ideation: Students were asked “During the past 12 months, did you ever seriously consider attempting suicide?”
5. Academic risk: Risk of academic failure including usually getting low grades and grades worse than others, and low commitment to school including school not meaningful or important for future, and cut school.

**Data Source:** Washington State Healthy Youth Survey 2016.



## Adult (Age 18+) Health Risk Behaviors




### Indicator Notes

1. Adult smoking: Respondents are asked “Have you smoked at least 100 cigarettes in your lifetime?” and “Do you still smoke?”
2. Adult marijuana: Respondents were asked “During the past 30 days, on how many days did you use marijuana or hashish?”
3. Binge drinking: Past 30 days, adult men having five or more drinks or adult women having four or more drinks on one occasion.
4. CDC recommends 150 minutes of moderate aerobic physical activity or 75 minutes of vigorous aerobic physical activity a week, combined with some form of muscle strengthening activity three times a week. People whose work involves mostly walking meet the aerobic recommendation. People whose work involves heavy labor meet both the strength and aerobic recommendations.
5. Nutrition: Respondents are asked a series of questions about fruits and vegetables eaten in the past month. CDC recommends three servings of vegetables and two servings of fruit a day. Very poor nutrition is defined here as eating fruits and vegetables less than once a day.
6. E-Cigarettes: Respondent is asked “During the past 30 days, on how many days did you use electronic cigarettes, also called E-cigarettes or vape pens?”

**Data Source:** Washington Behavioral Risk Factor Surveillance System 2013-2015

## Adult (Age 18+) Health Risk Conditions




### Indicator Notes

1. Obesity in adults is defined as body mass index  $\geq 30$  kg /m<sup>2</sup> based on self reported height and weight.
2. Self reported lifetime prevalence – Survey respondent answered “yes” to “have you ever been told by a healthcare professional that you have high blood pressure / high cholesterol?”
3. Food Insecurity: Respondents were asked “How often in the past 12 months would you say you were worried or stressed about having enough money to buy nutritious meals?” Responses of “sometimes”, “usually”, or “always” were considered to be food insecure.

**Data Source:** Washington Behavioral Risk Factor Surveillance System 2013-2015

## Adult (Age 18+) Preventive Care




### Indicator Notes

1. The Department of Health recommends women age 50 or older should have a mammogram every two years.
2. Flu vaccine: Respondent has not had a flu vaccine in the past year.
3. Personal physician: Respondent is asked: "Do you have one person you think of as your personal doctor or health care provider?"
4. Respondent reports needing to see a doctor, but could not due to cost in the past year.
5. No dental visit: Respondent reports it has been more than a year since they visited a dentist for any reason.
6. No checkup: Respondent reports it has been more than a year since they had a routine medical checkup.

**Data Source:** Washington Behavioral Risk Factor Surveillance System 2013-2015

## Adult (Age 18+) Chronic Disease




### Indicator Notes

1. Self reported lifetime prevalence – Survey respondent answered “yes” to “have you ever been told by a healthcare professional that you have asthma / diabetes / heart attack, coronary heart disease, or angina / arthritis / cancer?”

**Data Source:** Washington Behavioral Risk Factor Surveillance System 2013-2015.

## Adult (Age 18+) Quality of Life




### Indicator Notes

1. General health: respondent reports, in general, health is fair or poor.
2. Poor physical health: Respondent reports that on 14 or more of the past 30 days, their physical health was not good.
3. Poor mental health: Respondent reports that on 14 or more of the past 30 days, their mental health was not good.
4. Need medical equipment: Respondents are asked “Do you now have any health problem that requires you to use special equipment, such as a cane, a wheelchair, a special bed, or a special telephone?”
5. Activity limitation: Respondent is asked “Are you limited in any way in any activities because of physical, mental, or emotional problems?”

**Data Source:** Washington Behavioral Risk Factor Surveillance System 2013-2015

## Senior (Age 65+) Health Risks




### Indicator Notes


1. Living with chronic disease: Respondent is asked have you ever been told by a doctor or health care professional that you have ... arthritis / asthma / COPD / cancer / diabetes, heart disease / stroke / kidney disease.
2. Activities of daily living: Respondent is asked if they have serious difficulty ...seeing even with glasses / concentrating remembering or making decisions / walking or climbing stairs / dressing or bathing / doing errands alone such as visiting a doctor or shopping

**Data Source:** Washington Behavioral Risk Factor Surveillance System 2013-2015

# Mortality



**Cause of Death in Washington State and Non-Hispanic White Age standardized percent of all deaths.**



**Data Source:** Washington State Department of Health, Center for Health Statistics, Death Certificate Data, 2011-2015.

## Appendix: Data Sources & Definitions

The following provides references for more information on each data system and definitions of technical terms used in this report. Analyses for this report were completed using Stata/IC 13.0. Some estimates were obtained from previously published reports.

### **DATA SYSTEMS:**

#### **Office of Financial Management (OFM) Population Estimates**

- For more information on OFM intercensal population estimates, go to: <http://www.ofm.wa.gov/pop/default.asp>

#### **American Community Survey (ACS) and Public Use Microdata Sample (PUMS)**

- For more information on the American Community Survey, go to: <http://factfinder2.census.gov/faces/nav/jsf/pages/index.xhtml>
- For more information on Public Use Microdata Sample go to: [http://www.census.gov/acs/www/data\\_documentation/public\\_use\\_microdata\\_sample/](http://www.census.gov/acs/www/data_documentation/public_use_microdata_sample/)

#### **Pregnancy Risk Assessment Monitoring Survey (PRAMS)**

- For more information on PRAMS, go to: <http://www.doh.wa.gov/DataandStatisticalReports/HealthBehaviors/PregnancyRiskAssessmentMonitoringSystem.aspx>

#### **Washington Birth Certificate Data**

- For more information on birth data, go to: <http://www.doh.wa.gov/DataandStatisticalReports/VitalStatisticsData/BirthData.aspx>

#### **Washington State Department of Health, Office of Immunization and Child Profile**

- For more information on immunization data, go to: <http://www.doh.wa.gov/DataandStatisticalReports/SchoolImmunization.aspx>

#### **Washington State Healthy Youth Survey (HYS)**

- For more information on the HYS, go to: <http://www.doh.wa.gov/DataandStatisticalReports/HealthBehaviors/HealthyYouthSurvey.aspx> or <http://www.askhys.net/>
- For technical notes on the HYS, go to: <http://www.doh.wa.gov/DataandStatisticalReports/HealthBehaviors/HealthyYouthSurvey/TechnicalNotes.aspx>

#### **Behavioral Risk Factor Surveillance System (BRFSS)**

- For more information on Washington State BRFSS, go to: <http://www.doh.wa.gov/DataandStatisticalReports/HealthBehaviors/BehavioralRiskFactorSurveillanceSystemBRFSS.aspx>
- For more information on national BRFSS, go to: <http://www.cdc.gov/brfss>.

#### **Washington State Death Certificate Data**

- For more information on death records, go to: <http://www.doh.wa.gov/DataandStatisticalReports/VitalStatisticsData/DeathData.aspx>

#### **Washington State Cancer Registry (WSCR)**


- For more information on WSCR, go to: <https://fortress.wa.gov/doh/wscr/WSCR/>



**CONFIDENCE INTERVALS:**

Most of the estimates provided in this report come with some intrinsic level of uncertainty due to the random nature of the data. Statistical uncertainty can be summarized by a 95% confidence interval, also called the margin of error. 95% confidence means that, if the survey were repeated in exactly the same way with a different random sample of people, the new estimate would fall within the confidence interval 95% of the time. Confidence intervals are represented on graphs by whisker bars above and below the estimate.

**Interpreting Margin of Error**



**UNRELIABLE DATA:**

Estimates based on too few respondents are considered to be unreliable, and may constitute a breach of confidentiality in some circumstances. In this report data with a numerator < 10, or a denominator < 50, or a relative standard error > 30% are not reported.

**AGE-ADJUSTED PERCENT:**

Percentages that have been adjusted to control for differences in age when comparing two demographic groups. Respondents are weighted to match the US Census 2000 standard population.

## **GLOSSARY:**

---

<sup>i</sup> Prevalence: The fraction of the population with a condition at a particular point in time, typically expressed as a percent.

<sup>ii</sup> Life course approach: A philosophy of public health that recognizes the importance of promoting health at all life stages.

<sup>iii</sup> Epigenetic: Conditions in the mother prior to conception can affect how certain genes are expressed in the child.